


UTILITY PATENT APPLICATION TRANSMITTAL <small>(Only for new nonprovisional applications under 37 CFR 1.53(b))</small>	Attorney Docket No.	0010-1066-0
	First Inventor or Application Identifier	Vitaliy A. LIVSHITS, et al.
	Title	METHOD FOR PRODUCING L-AMINO ACID



APPLICATION ELEMENTS <i>See MPEP chapter 600 concerning utility patent application contents</i>	ADDRESS TO: Assistant Commissioner for Patents Box Patent Application Washington, DC 20231
<p>1. <input checked="" type="checkbox"/> Fee Transmittal Form (e.g. PTO/SB/17) (Submit an original and a duplicate for fee processing)</p> <p>2. <input checked="" type="checkbox"/> Specification Total Pages 58</p> <p>3. <input type="checkbox"/> Drawing(s) (35 U.S.C. 113) Total Sheets </p> <p>4. <input checked="" type="checkbox"/> Oath or Declaration Total Pages 4</p> <p>a. <input checked="" type="checkbox"/> Newly executed (original)</p> <p>b. <input type="checkbox"/> Copy from a prior application (37 C.F.R. §1.63(d)) (for continuation/divisional with box 15 completed)</p> <p>i. <input type="checkbox"/> </p> <p>DELETION OF INVENTOR(S) Signed statement attached deleting inventor(s) named in the prior application, see 37 C.F.R. §1.63(d)(2) and 1.33(b).</p> <p>5. <input type="checkbox"/> Incorporation By Reference (usable if box 4B is checked) The entire disclosure of the prior application, from which a copy of the oath or declaration is supplied under Box 4B, is considered to be part of the disclosure of the accompanying application and is hereby incorporated by reference therein.</p>	ACCOMPANYING APPLICATION PARTS <p>6. <input type="checkbox"/> Assignment Papers (cover sheet & document(s))</p> <p>7. <input type="checkbox"/> 37 C.F.R. §3.73(b) Statement <input type="checkbox"/> Power of Attorney (when there is an assignee)</p> <p>8. <input type="checkbox"/> English Translation Document (if applicable)</p> <p>9. <input type="checkbox"/> Information Disclosure Statement (IDS)/PTO-1449 <input type="checkbox"/> Copies of IDS Citations</p> <p>10. <input type="checkbox"/> Preliminary Amendment</p> <p>11. <input checked="" type="checkbox"/> White Advance Serial No. Postcard</p> <p>12. <input type="checkbox"/> Small Entity Statement(s) <input type="checkbox"/> Statement filed in prior application. Status still proper and desired.</p> <p>13. <input checked="" type="checkbox"/> Certified Copy of Priority Document(s)(2) (if foreign priority is claimed)</p> <p>14. <input checked="" type="checkbox"/> Other:</p> <p>Notice of Priority, Receipt of an Original Deposit of Microorganisms for the Purposes of Patent Procedure (VKPM B-7707, VKPM B-7708, VKPM B-7712, VKPM B-7713, VKPM B-7714, VKPM B-7715, VKPM B-7716, VKPM B-7718, VKPM B-7719, VKPM B-7722, VKPM B-7728, VKPM B-7729, VKPM B-7730, VKPM B-7731, VKPM B-7748, VKPM B-7752, VKPM B-7753, VKPM B-7754)</p>
<p>15. If a CONTINUING APPLICATION, check appropriate box, and supply the requisite information below:</p> <p><input type="checkbox"/> Continuation <input type="checkbox"/> Divisional <input type="checkbox"/> Continuation-in-part (CIP) of prior application no.:</p> <p>Prior application information: Examiner: Group Art Unit:</p>	
<p>16. Amend the specification by inserting before the first line the sentence:</p> <p><input type="checkbox"/> This application is a <input type="checkbox"/> Continuation <input type="checkbox"/> Division <input type="checkbox"/> Continuation-in-part (CIP) of application Serial No. Filed on</p> <p><input type="checkbox"/> This application claims priority of provisional application Serial No. Filed</p>	
<p>17. CORRESPONDENCE ADDRESS</p> <p>OBLON, SPIVAK, MCCLELLAND, MAIER & NEUSTADT, P.C. FOURTH FLOOR 1755 JEFFERSON DAVIS HIGHWAY ARLINGTON, VIRGINIA 22202 (703) 413-3000 FACSIMILE: (703) 413-2220</p>	

Name:	Norman F. Oblon	Registration No.:	24,618
Signature:	MARVIN J. SPIVAK REGISTRATION NUMBER 24,913	Date:	12/13/99
Name:		Registration No.:	

METHOD FOR PRODUCING L-AMINO ACID

Technical Field

5 The present invention relates to a method for producing an amino acid. In particular, the present invention relates to an L-amino acid-producing bacterium belonging to the genus *Escherichia* and a method for producing L-amino acids, more specifically, L-glutamic acid, L-lysine, L-threonine, 10 L-alanine, L-histidine, L-proline, L-arginine, L-valine, and L-isoleucine, using the bacterium.

Background Art

15 For production of an L-amino acid by fermentation, a strain isolated from the natural world or an artificial mutant of the strain has been used to improve productivity. For example, in the case of L-lysine, many artificial mutants producing L-lysine are known, and most of them are mutants resistant to S-2-aminoethylcysteine (AEC) and belong to the 20 genus *Brevibacterium*, *Corynebacterium*, *Bacillus* or *Escherichia*. Also, there have been proposed various technics for increasing amino acid production such as use of a transformant obtained by using a recombinant DNA (U.S. Patent No. 4,278,765).

25 The technics are mostly based on enhancement of an activity of an enzyme involved in an amino acid biosynthetic pathway, conversion of the enzyme to that desensitized in

inhibition and the like (As to bacterium belonging the genus *Escherichia*, see Japanese Patent Application Laid-Open No. 56-18596 (1981) and International Publication No. WO 95/16042).

5 On the other hand, as an example of improvement of amino acid productivity by enhancing an amino acid excretion protein, a bacterium belonging to the genus *Corynebacterium* in which an L-lysine excretion gene, *lyse* is enhanced is known. However, as to bacteria belonging to the genus
10 *Escherichia*, it is unknown even whether an L-amino acid excretion protein is present or not. Therefore, it is unknown whether enhancement of the L-amino acid excretion protein is effective in L-amino acid production using a bacterium belonging to the genus *Escherichia* or not.

15 Although the entire nucleotide sequence of *E. coli* strain K-12 belonging to the genus *Escherichia* has been already determined (Science, 277, 1453-1474(1997)), there are a large number of proteins of which functions are unknown.

20

Disclosure of the Invention

An object of the present invention is to obtain a protein participating in excretion of an L-amino acid, thereby providing a strain improved in L-amino acid
25 productivity and an improved method for producing an L-amino acid by fermentation.

The inventors have conducted screening for the protein

participating in excretion of an L-amino acid. As a result, the present inventors have found that a yield of an L-amino acid based on consumed sugar is increased when a particular gene is enhanced. On the basis of the finding, the present invention has been completed.

Thus, the present invention provides a bacterium belonging to the genus *Escherichia* and having an ability to produce an L-amino acid, wherein the ability to produce the L-amino acid is increased by increasing an expression amount of at least one protein selected from the group consisting of the following proteins of (A) to (H) (hereinafter also referred to as "the bacterium of the present invention"):

(A) a protein having an amino acid sequence shown in SEQ ID NO: 10 in Sequence Listing;

(B) a protein which has an amino acid sequence including deletion, substitution, insertion, addition or inversion of one or several amino acids in the amino acid sequence shown in SEQ ID NO: 10 in Sequence Listing, and which has an activity of increasing the ability to produce the L-amino acid of the bacterium having the protein;

(C) a protein having an amino acid sequence shown in SEQ ID NO: 12 in Sequence Listing;

(D) a protein which has an amino acid sequence including deletion, substitution, insertion, addition or inversion of one or several amino acids in the amino acid sequence shown in SEQ ID NO: 12 in Sequence Listing, and which has an activity of increasing the ability to produce the

L-amino acid of the bacterium having the protein;

(E) a protein having an amino acid sequence shown in SEQ ID NO: 14 in Sequence Listing;

(F) a protein which has an amino acid sequence
5 including deletion, substitution, insertion, addition or inversion of one or several amino acids in the amino acid sequence shown in SEQ ID NO: 14 in Sequence Listing, and which has an activity of increasing the ability to produce the L-amino acid of the bacterium having the protein;

10 (G) a protein having an amino acid sequence shown in SEQ ID NO: 16 in Sequence Listing; or

(H) a protein which has an amino acid sequence
including deletion, substitution, insertion, addition or inversion of one or several amino acids in the amino acid
15 sequence shown in SEQ ID NO: 16 in Sequence Listing, and which has an activity of increasing the ability to produce the L-amino acid of the bacterium having the protein.

The bacterium of the present invention preferably an L-lysine-producing bacterium in which an expression amount
20 of at least one protein selected from the group consisting of the proteins (A) to (D), (G) and (H) is increased; an L-glutamic acid-producing bacterium in which an expression amount of at least one protein selected from the group consisting of the proteins (A) to (H) is increased; an
25 L-alanine-producing bacterium in which an expression amount of at least one protein selected from the group consisting of the proteins (C) and (D) is increased; an L-valine-

producing bacterium in which an expression amount of at least one protein selected from the group consisting of the proteins (C) and (D) is increased; an L-histidine-producing bacterium in which an expression amount of at least one protein selected from the group consisting of said proteins (C) to (F) is increased; an L-proline-producing bacterium in which an expression amount of at least one protein selected from the group consisting of said proteins (A) to (F) is increased; an L-threonine-producing bacterium in which an expression amount of at least one protein selected from the group consisting of said proteins (E) and (F) is increased; an L-arginine-producing bacterium in which an expression amount of at least one protein selected from the group consisting of said proteins (G) and (H) is increased; or an L-isoleucine-producing bacterium in which an expression amount of at least one protein selected from the group consisting of said proteins (C) and (D) is increased.

Preferably, in the bacterium of the present invention, a copy number of a DNA coding for said protein in a cell is increased. The DNA is preferably carried on a multicopy vector in the cell or on a transposon in the cell.

The present invention also provides a method for producing an L-amino acid, comprising the steps of:

cultivating the bacterium of the present invention in a culture medium, to produce and accumulate the L-amino acid in the medium, and

recovering the L-amino acid from the medium

(hereinafter also referred to as "the bacterium of the present invention").

The method of the present invention preferably an L-lysine production method using an L-lysine-producing bacterium in which an expression amount of at least one protein selected from the group consisting of the proteins (A) to (D), (G) and (H) is increased; an L-glutamic acid production method using an L-glutamic acid-producing bacterium in which an expression amount of at least one protein selected from the group consisting of the proteins (A) to (H) is increased; an L-alanine production method using an L-alanine-producing bacterium in which an expression amount of at least one protein selected from the group consisting of the proteins (C) and (D) is increased; an L-valine production method using an L-valine-producing bacterium in which an expression amount of at least one protein selected from the group consisting of the proteins (C) and (D) is increased; an L-histidine production method using an L-histidine-producing bacterium in which an expression amount of at least one protein selected from the group consisting of said proteins (C) to (F) is increased; an L-proline production method using an L-proline-producing bacterium in which an expression amount of at least one protein selected from the group consisting of said proteins (A) to (F) is increased; an L-threonine production method using an L-threonine-producing bacterium in which an expression amount of at least one protein selected from the

group consisting of said proteins (E) and (F) is increased;
an L-arginine production method using an L-arginine-
producing bacterium in which an expression amount of at least
one protein selected from the group consisting of said
5 proteins (G) and (H) is increased; or an L-isoleucine
production method using an L-isoleucine-producing bacterium
in which an expression amount of at least one protein
selected from the group consisting of said proteins (C) and
(D) is increased.

10 Preferably, in the method of the present invention, a
copy number of a DNA coding for said protein in a cell of
the bacterium is increased. The DNA is preferably carried
on a multicopy vector in the cell, or on a transposon in the
cell.

15 According to the present invention, an ability to
produce an L-amino acid of a bacterium belonging to the genus
Escherichia can be increased. Also, a method for producing
an L-amino acid can be improved in a production rate of an
L-amino acid.

20 The present invention will be explained in detail below.
Hereinafter, an amino acid is of L-configuration unless
otherwise noted.

<1> Bacterium of the present invention

The bacterium of the present invention is a bacterium
25 belonging to the genus *Escherichia* and having an ability to
produce an amino acid, in which the ability to produce the
amino acid is increased by increasing an expression amount

of a protein which has an activity of increasing the ability to produce the amino acid of the bacterium, or an activity of increasing resistance to an amino acid or amino acid analogue. Hereinafter, the protein is referred to as "amino acid excretion protein" for the sake of convenience.

However, the term does not mean that function of the protein is limited to amino acid excretion.

Examples of the amino acid excretion protein include a protein having an amino acid sequence shown in SEQ ID NO: 10, a protein having an amino acid sequence shown in SEQ ID NO: 12, a protein having an amino acid sequence shown in SEQ ID NO: 14 and a protein having an amino acid sequence shown in SEQ ID NO: 16.

The amino acid excretion protein may have selectivity to amino acid. An amino acid excretion protein appropriate for each amino acid can be determined by allowing the amino acid excretion protein to be expressed in a bacterium belonging to the genus *Escherichia* and having an ability to produce the amino acid, and measuring an increase of a yield of the amino acid or measuring an increase of a minimum inhibition concentration (MIC) of an amino acid or amino acid analogue.

For example, in the case of lysine, a protein having an amino acid sequence shown in SEQ ID NO: 10, 12 or 16 is effective; in the case of glutamic acid, a protein having an amino acid sequence shown in SEQ ID NO: 10, 12, 14 or 16 is effective; in the case of alanine, a protein having an

amino acid sequence shown in SEQ ID NO: 12 is effective; in the case of valine, a protein having an amino acid sequence shown in SEQ ID NO: 12 is effective; in the case of histidine, a protein having an amino acid sequence shown in SEQ ID NO: 12 or 14; in the case of proline, a protein having an amino acid sequence shown in SEQ ID NO: 10, 12 or 14 is effective; in the case of threonine, a protein having an amino acid sequence shown in SEQ ID NO: 14 is effective; in the case of arginine, a protein having an amino acid sequence shown in SEQ ID NO: 16 is effective; and in the case of isoleucine, a protein having an amino acid sequence shown in SEQ ID NO: 12 is effective.

The term "an expression amount is increased" used herein usually means that the expression amount is larger than that in a wild strain of *E. coli* such as strain MG1655 or W3110. The terms also means that when a strain is obtained by modification through genetic engineering technics or the like, the expression amount is larger than that prior to the modification. The expression amount of the amino acid excretion protein may be determined directly by the determination of the amino acid excretion protein or indirectly by the determination of MIC of an amino acid or amino acid analogue or of amino acid productivity of a bacterium belonging to the genus *Escherichia* and having the amino acid excretion protein.

The method for increasing the expression amount of the amino acid excretion protein is exemplified by a method for

increasing a copy number of DNA encoding the amino acid excretion protein in a cell of the bacterium.

For increasing the copy number in the cell, a DNA fragment coding for the amino acid excretion protein may be
5 ligated to a vector which functions in a bacterium belonging to the genus *Escherichia* to produce a recombinant DNA, which is introduced to a host to transform it. The copy number of the gene coding for the amino acid excretion protein (amino acid excretion protein gene) in the cell of the
10 transformant strain increases, thereby increasing the expression amount of the amino acid excretion protein. The vector is preferably a multicopy vector.

The increase of the copy number in the cell can be achieved by allowing plural copies of the amino acid
15 excretion protein gene to exist on chromosomal DNA of the host. The introduction of plural copies of the amino acid excretion protein gene to chromosomal DNA of a bacterium belonging to the genus *Escherichia*, may be conducted through homologous recombination by using a sequence of which plural
20 copies exist on the chromosomal DNA, as a target. As the sequence of which plural copies exist on the chromosomal DNA, a repetitive DNA and an inverted repeat present in a terminal portion of a transposable element may be used.

Alternatively, as disclosed in Japanese Patent Application
25 Laid-Open No. 2-109985 (1990), the plural copies can be introduced to the chromosomal DNA by making the amino acid excretion protein gene carried on a transposon and allowing

the transposon to be transposed, which is preferred. According to any of the above-mentioned methods, the copy number of the amino acid excretion protein gene in the transformant strain increases, thereby increasing the expression amount of the amino acid excretion protein.

The multicopy vector is exemplified by plasmid vectors such as pBR322, pMW118, pUC19 or the like, and phage vectors such as λ 1059, λ BF101, M13mp9 or the like. The transposon is exemplified by Mu, Tn10, Tn5 or the like.

The introduction of a DNA into a bacterium belonging to the genus *Escherichia* can be performed, for example, by a method of D. M. Morrison (Methods in Enzymology 68, 326 (1979)) or a method in which recipient bacterial cells are treated with calcium chloride to increase permeability of DNA (Mandel, M. and Higa, A., J. Mol. Biol., 53, 159 (1970)) and the like.

Besides the above-mentioned gene amplification, the increase of the expression amount of the amino acid excretion protein can be also achieved by replacing an expression regulatory sequence such as a promoter of the amino acid excretion protein gene with stronger one (see Japanese Patent Application Laid-Open No. 1-215280 (1989)). For example, *lac* promoter, *trp* promoter, *tac* promoter, P_R promoter and P_L promoter of lambda phage, and the like are known as a strong promoter. The replacement with the promoter enhances expression of the amino acid excretion protein, thereby increasing the expression amount of the

amino acid excretion protein. The enhancement of the expression regulatory sequence may be combined with the increase of the copy number of the amino acid excretion protein.

5 In the bacterium of the present invention, expression amounts of plural amino acid excretion proteins may be increased.

The amino acid excretion protein is encoded by genes which are known as *yahN* gene, *yeaS* gene, *yfiK* gene and *yggA* gene and of which functions are unknown. Therefore, the DNA
10 encoding the amino acid excretion protein can be obtained by synthesizing primers based on the known sequences (for example, the entire nucleotide sequence of chromosome of *Escherichia coli* strain K-12 has been already determined
15 (Science, 277, 1453-1474(1997))), and conducting amplification by PCR using chromosomal DNA of a bacterium belonging to the genus *Escherichia* as a template. Also, the object DNA fragment can be selected by hybridization from a chromosomal DNA library of a bacterium belonging to the
20 genus *Escherichia* by preparing a probe based on the known sequences. Alternatively, the DNA encoding the amino acid excretion protein may be synthesized based on the known sequences. The nucleotide sequence of the DNA encoding the amino acid excretion protein is exemplified by that shown
25 in SEQ ID NO: 9, 11, 13 or 15 in the Sequence Listing.

Methods for preparation of chromosomal DNA, preparation of chromosomal DNA library, hybridization, PCR,

preparation of plasmid DNA, digestion and ligation of DNA, transformation, selection of an oligonucleotide as a primer and the like may be ordinary methods well known to one skilled in the art. These methods are described in Sambrook, J., Fritsch, E. F., and Maniatis, T., "Molecular Cloning A Laboratory Manual, Second Edition", Cold Spring Harbor Laboratory Press (1989) and the like.

The amino acid excretion protein may comprise substitution, deletion, insertion, addition or inversion of one or several amino acids at one or a plurality of positions, provided that the activity of increasing the ability to produce the amino acid of the bacterium belonging to the genus *Escherichia* and having the protein is not deteriorated. The term "several" may vary depending on a position in a steric structure of the protein and a kind of an amino acid residue. It is because some amino acids such as isoleucine and valine have high similarity to each other, and a difference between such the amino acids does not largely affect the steric structure of the protein.

The DNA which codes for the substantially same protein as the amino acid excretion protein as described above, may be obtained, for example, by modifying the nucleotide sequence, for example, by means of the site-directed mutagenesis method so that one or more amino acid residues at a specified site involve substitution, deletion, insertion, addition or inversion. The DNA modified as described above may be obtained by the conventionally known

mutation treatment. The mutation treatment includes a method for treating a DNA coding for the amino acid excretion protein *in vitro*, for example, with hydroxylamine, and a method for treating a microorganism, for example, a bacterium belonging to the genus *Escherichia*, harboring a DNA coding for the amino acid excretion protein with ultraviolet irradiation or a mutating agent such as N-methyl-N'-nitro-N-nitrosoguanidine (NG) and nitrous acid usually used for the mutation treatment.

The substitution, deletion, insertion, addition or inversion of the one or more amino acid residues includes a naturally-occurring mutation or variation which is resulted from a difference between individual microorganisms having the amino acid excretion protein and a difference between species, strains or the like.

The DNA, which codes for substantially the same protein as the amino acid excretion protein, can be obtained by allowing a DNA having the mutation as described above to be expressed in a cell of an appropriate bacterium belonging to the genus *Escherichia*, and investigating the increase of amino acid productivity of the cell.

Also, the DNA, which codes for substantially the same protein as the amino acid excretion protein, can be obtained by isolating a DNA which hybridizes with DNA having, for example, a nucleotide sequence shown in SEQ ID NO: 9, 11, 13 or 15 in Sequence Listing under stringent conditions, and which codes for a protein having the activity of increasing

the ability to produce the amino acid of the bacterium belonging to the genus *Escherichia*, from DNAs encoding the amino acid excretion proteins having mutations or cells containing the DNAs. The term "stringent conditions" referred to herein means a condition under which a specific hybrid is formed, and a non-specific hybrid is not formed. It is difficult to clearly express this condition by using any numerical value. However, for example, the stringent conditions include a condition under which DNAs having high homology, for example, DNAs having homology of not less than 70% with each other are hybridized, and DNAs having homology lower than the above with each other are not hybridized, or a condition of a salt concentration corresponding to 60°C, 1x SSC, 0.1% SDS, preferably 0.1x SSC, 0.1% SDS which is a washing condition of ordinary Southern hybridization.

Although there may be a gene in which a stop codon is made in the middle, or a gene encoding a protein losing the activity due to mutation of the active center among the genes which hybridize under such the condition, such genes can be easily eliminated by ligating the genes to a commercially available activity-expression vector and determining the activity of increasing the ability to produce the amino acid of the bacterium belonging to the genus *Escherichia* as described above.

The term "DNA coding for a protein" used herein means a DNA of which one of strands codes for the protein when the DNA is double-stranded.

By increasing an expression amount of an amino acid excretion protein in an amino acid-producing bacterium belonging to the genus *Escherichia* as described above, a produced amount of the amino acid can be increased. As the bacterium belonging to the genus *Escherichia* in which the expression amount of the amino acid excretion protein is to be increased, strains which have abilities to produce desired amino acids (amino acid productivities) are used. Besides, an ability to produce an amino acid may be imparted to a bacterium in which the expression amount of the amino acid excretion protein is increased. Examples of amino acid-producing bacteria belonging to the genus *Escherichia* include *E. coli* AJ13199 (FR patent No. 2747689), and those obtainable from known materials (e.g., *E. coli* W3110 (tyrA)/pCABD2, *E. coli* VL614, *E. coli* VL2054, *E. coli* VL2160, *E. coli* VL2151, *E. coli* W3350 *argE*::Tn10/pKA10 as described in the Examples below).

For reference, the amino acid excretion protein according to the present invention was identified for the first time as described below.

The present inventors have identified *rhtB* and *rhtC* as threonine excretion protein genes of a bacterium belonging to the genus *Escherichia*. The present inventors searched databases based on a hypothesis that amino acid excretion proteins may share a common structure. Namely, BLAST and PSI-BLAST search (Altschul, S.F. et al., Nucleic Acids Res., 25, 3389-3402(1997)) for homology of a protein encoded by

rhtB was performed in GenBank CDS, PDB, SWISS-PROT, Spupdate and PIR. Tblastn search was performed in unfinished microbial genomes. BLITZ search (Sturrock, S.S., and Collins, J.F., Mpsch version 1.3. Biocomputing research unit University of Edinburgh, UK (1993)) was performed in SWALL database. SMART search (Ogiwara, I. et al., Protein Sci., 5, 1991-1999 (1996)) was performed in the databases of translations and SWISS-PROT. From the samples of more than 60 sequences found, YeaS (corresponding to f212 of ACCESSION No. AE000274 in GenBank), YahN (corresponding to f223 of ACCESSION No. AE000140 in GenBank), YfiK (corresponding to o195 of ACCESSION No. AE000344 in GenBank) and YggA (corresponding to f211 of ACCESSION No. AE000375 in GenBank) remained as proteins which may have similar function to RhtB, among those originating from *E. coli*. Since functions of any of these genes were unknown, the genes were actually obtained, and effects thereof on MIC of amino acids and amino acid analogues and on amino acid production were examined by enhancing activities thereof. As a result, an effect of increasing MIC of some amino acids and analogues was found with respect to YeaS, YfiK, YahN and YggA. Further examination has revealed that proteins encoded by these genes exhibit an effect of increasing an amino acid accumulation, although they may have some amino acid selectivities.

<2> Method of the present invention

The method of the present invention comprises the steps of cultivating the bacterium of the present invention, in a culture medium, to produce and accumulate the amino acid in the medium, and recovering the amino acid from the medium.

5 Suitable amino acids include lysine, glutamic acid, alanine, valine, homoserine, proline, and threonine.

In the method of present invention, the cultivation of the bacterium belonging to the genus *Escherichia*, the collection and purification of amino acid from the liquid
10 medium may be performed in a manner similar to those of the conventional method for producing an amino acid by fermentation using a bacterium. A medium used in cultivation may be either a synthetic medium or a natural medium, so long as the medium includes a carbon and a nitrogen
15 source and minerals and, if necessary, nutrients which the bacterium used requires for growth in appropriate amounts. The carbon source may include various carbohydrates such as glucose and sucrose, and various organic acids. Depending on assimilatory ability of the used bacterium, alcohol
20 including ethanol and glycerol may be used. As the nitrogen source, ammonia, various ammonium salts such as ammonium sulfate, other nitrogen compounds such as amines, a natural nitrogen source such as peptone, soybean hydrolyte and digested fermentative microbe are used. As minerals,
25 monopotassium phosphate, magnesium sulfate, sodium chloride, ferrous sulfate, manganese sulfate, calcium carbonate are used.

The cultivation is preferably culture under an aerobic condition such as a shaking culture, and an aeration and stirring culture. The temperature of culture is usually 20 to 40°C, preferably 30 to 38°C. The pH of the culture is usually between 5 and 9, preferably between 6.5 and 7.2. The pH of the culture can be adjusted with ammonia, calcium carbonate, various acids, various bases, and buffers. Usually, a 1 to 3-day cultivation leads to the accumulation of the target amino acid in the medium.

Recovering the amino acid can be performed by removing solids such as cells from the medium by centrifugation or membrane filtration after cultivation, and then collecting and purifying the target amino acid by ion exchange, concentration and crystalline fraction methods and the like.

Best Mode for Carrying Out the Invention

The present invention will be more concretely explained below with reference to Examples.

Example 1. Preparation of the DNA fragments which code for amino acid excretion proteins.

The entire nucleotide sequence of chromosome of *E. coli* strain K-12 has been determined (Science, 277, 1453-1474, 1997). Based on the reported nucleotide sequence, primers were synthesized and the genes *yahN*, *yfiK*, *yeaS* and *yggA* were amplified by PCR.

(1). Chromosomal DNA of the *E. coli* strain MG1655 was used as a template.

The chromosomal DNA was prepared by an ordinary method (Sambrook, J., Fritsch E. F. and Maniatis T. (1989) Molecular cloning: a laboratory manual, 2nd ed. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N. Y.). In the PCR reaction, a standard condition described in "PCR protocols. Current methods and applications". (White, B.A., ed. Humana Press, Totowa, New Jersey, 1993) was used. The obtained PCR products were purified by an ordinary method and digested with restriction enzymes as described below.

The *yahN* gene was amplified by using the primers No.1 and No. 2.

Primer No.1: gtgtggaaccgacgccgat (a sequence complementary to a sequence of from 1885 base to 1904 base in a nucleotide sequence registered under ACCESSION No. AE000140 in GenBank; SEQ ID NO: 17), and

Primer No.2: tgttgtatggtacgggggttcgag (a sequence of from 223 base to 245 base in the same; SEQ ID NO: 18).

The obtained PCR product after purification was digested with restriction enzymes *Pst*I and *Stu*I and ligated to vector pUC21 (Vieira, Messing, Gene, 100, 189-194, 1991) digested with the enzymes *Pst*I and *Eco*RV by using a ligation kit. Then, transformation of competent cells of *E. coli* TG1 (Sambrook, J., Fritsch E. F. and Maniatis T. (1989) Molecular cloning: a laboratory manual, 2nd ed. Cold Spring Harbor Laboratory

Press, Cold Spring Harbor, N. Y.) with the product was conducted and the cells were spread on L medium (10 g/l Bacto trypton, 5 g/l Yeast extract, 5 g/l NaCl, 15 g/l agar, pH 7.0) containing 10 mg/ml IPTG (isopropyl- β -D-thiogalactopyranoside) and 40 mg/ml X-gal (5-bromo-4-chloro-3-indolyl- β -D-galactoside) and 100 mg/ml ampicillin, and cultured overnight. Appeared white colonies were picked up and subjected to single colony isolation to obtain transformants. Plasmid was prepared from the transformants using an alkali extraction method and designated as pYAHN.

The *yeaS* gene was amplified by using the primers No.3 and No. 4.

Primer No.3: ctttgccaatcccgtctccc (a sequence complementary to a sequence of from 7683 base to 7702 base in a nucleotide sequence registered under ACCESSION No AE000274 in GenBank; SEQ ID NO: 19);

Primer No.4: gccccatgcataacggaaag (a sequence of from 5542 base to 5561 base in the same; SEQ ID NO: 19).

The obtained PCR product after purification was digested with a restriction enzyme *AvaI* and ligated to vector pUC19. After transformation of *E. coli* TG1 as above, the plasmid designated as pYEAS was obtained.

The *yfiK* gene was amplified by using the primers No.5 and No.6.

Primer No.5: gaagatcttgtaggccggataaggcg (a sequence of from 4155 base to 4177 base in a nucleotide sequence

registered under ACCESSION No AE000344 in GenBank, with a restriction enzyme *Bgl*III site added at the 5'-end thereof; SEQ ID NO: 21)

Primer No.6: tggttttaccaattggccgc (a sequence
5 complementary to a sequence of from 6307 base to 6326 base in the same; SEQ ID NO: 22).

The obtained PCR product after purification was digested with restriction enzymes *Bgl*III and *Mun*I and ligated to vector pUC21 digested with restriction enzymes *Bgl*III and *Eco*RI.
10 After transformation of *E. coli* TGI as above, the plasmid designated pYFIK was obtained.

The *yggA* gene was amplified by using the primers No.7 and No.8.

Primer No.7: acttctcccgcgagccagttc (a sequence
15 complementary to a sequence of from 9606 base to 9626 base in a nucleotide sequence registered under ACCESSION No AE000375 in GenBank; SEQ ID NO: 23).

Primer No.8: ggcaagcttagcgcctctgtt (a sequence of from 8478 base to 8498 base in the same; SEQ ID NO: 24).
20 The obtained PCR product after purification was digested with restriction enzymes *Hind*III and *Cla*I and ligated to vector pOK12 (Vieira, Messing, Gene, 100, 189-194, 1991) digested with the same restriction enzymes. After transformation of *E. coli* TGI as above, the plasmid
25 designated pYGGA was obtained.

(2). Chromosomal DNA of the *E. coli* strain W3110 was used as a template.

The *yahN* gene was amplified by using the primers No.9 (SEQ ID NO 1) and No. 10 (SEQ ID NO.2)

5 The *yeaS* gene was amplified by using the primers No.11 (SEQ ID NO 3) and No.12 (SEQ ID NO 4)

The *yfiK* gene was amplified by using the primers No.13 (SEQ ID NO 5) and No.14 (SEQ ID NO 6).

10 The *yggA* gene was amplified by using the primers No.15 (SEQ ID NO 7) and No.16 (SEQ ID NO 8)

The obtained PCR product was purified, digested with restriction enzymes *SacI* and *XbaI* (*EcoRI* and *PstI* for *yggA*), and ligated to plasmid pMW118 (Nippon Gene). The plasmid into which a DNA fragment of which sequence was identical to the reported sequence was inserted was designated as follows:

One carrying *yahN*: pMW118::*yahN*
 One carrying *yeaS*: pMW118::*yeaS*
 One carrying *yfiK*: pMW118::*yfiK*
 20 One carrying *yggA*: pMW118::*yggA*

Example 2. Effect of the *yahN*, *yeaS*, *yfiK*, and *yggA* DNA fragments amplification on the *E. coli* TGl resistance to some amino acids and amino acid analogues.

25 The homology of the *yeaS*, *yfiK*, *yahN* and *yggA* gene products with the lysine transporter, *Lyse*, of

Corynebacterium glutamicum (Vrljic et al., Mol. Microbiol., 22, 815-826, 1996) and RhtB protein involved in homoserine excretion, indicates the analogues function for these proteins. It is well known that the increased expression of the genes involved in antibiotic and heavy metal efflux increases the level of resistance to the drugs (Nikaido, H. J. Bacteriology, 178, 5853-5859, 1996). Therefore, the effect of the pYEAS, pYAHN, pYFIK, and pYGGA plasmids on susceptibility of the strain TG1 to some amino acids and amino acid analogues was tested. Overnight cultures of the *E. coli* strains TG1/pYEAS, TG1/pYAHN, TG1/pYFIK, TG1/pYGGA and of the control strains TG1/pUC21, TG1/pUC19 and TG1/pOK12 grown in M9 minimal medium with an appropriate antibiotic on a rotary shaker (10^9 cfu/ml) were diluted 1:100 in M9 minimal medium and grown for 5 h in the same medium. Then the log phase cultures thus obtained were diluted and about 10^4 alive cells were applied to well-dried test plates with M9 agar containing doubling increments of amino acids or analogues. Thus the minimum inhibition concentration (MIC) of these compounds were examined.

The results are shown in Table 1. It follows from the Table 1 that multiple copies of *yfiK* gene conferred increased resistance to proline, homoserine, histidine, threonine, glutamate, lysine, α -amino- β -hydroxyvaleric-acid (AHVA), S-(2-aminoethyl)-L-cysteine (AEC) and α -aminobutyric acid; multiple copies of *yahN* gene conferred increased resistance to proline, multiple copies of *yeaS* gene conferred increased

resistance to threonine, homoserine, lysine, glutamate, histidine, proline and α -aminobutyric acid; multiple copies of *yggA* gene conferred increased resistance to S-(2-aminoethyl)-L-cysteine (AEC), lysine, and arginine. These results indicate that except for *YahN*, every of the presumed transporters have specificity to several substrates (amino acids and amino acid analogues), or may show non-specific effects as a result of amplification.

Table 1

Substrate	MIC (μ g/ml) for <i>E. coli</i> TGI, harboring the plasmid				
	pUC21	pYFIK	pYAHN	pYEAS	pYGGA
L-homoserine	500	1000	500	1000	500
L-threonine	30000	40000	30000	50000	30000
L-lysine	5000	7500	5000	7500	15000
L-glutamate (Na salt)	5000	10000	5000	20000	5000
L-histidine	5000	10000	5000	30000	5000
L-valine	0.5	0.5	0.5	0.5	0.5
L-proline	1000	5000	2000	2000	1000
L-arginine	10000	10000	10000	10000	20000
AHVA	100	200	100	100	100
AEC	5	10	5	5	200
α -aminobutyric acid	2500	5000	2500	>10000	2500
4-aza-DL-leucine	100	100	100	100	100

Example 3. Effect of *yeaS*, *yahN*, and *yfiK* DNA fragments amplification on glutamic acid production.

The *E. coli* strain AJ13199 (FR patent No. 2747689) was transformed with the vector pUC21 and each of the plasmids

pYAHN, pYEAS and pYFIK. Thus the strains AJ13199/pUC21 (VKPM B-7728), AJ13199/pYAHN (VKPM B-7729), AJ13199/pYEAS (VKPM B-7731), and AJ13199/pYFIK (VKPM B-7730) were obtained.

5 These strains were each cultivated at 37°C for 18 hours in a nutrient broth with 100 mg/l ampicillin, and 0.3 ml of the obtained culture was inoculated into 3 ml of a fermentation medium containing 100 mg/l ampicillin, in a 20 x 200 mm test tube, and cultivated at 37°C for 48 hours with
10 a rotary shaker. After the cultivation, an accumulated amount of glutamic acid in the medium was determined by known method.

The composition of the fermentation medium (g/l):

15	Glucose	80
	(NH ₄) ₂ SO ₄	22
	K ₂ HPO ₄	2
	NaCl	0.8
	MgSO ₄ · 7H ₂ O	0.8
20	FeSO ₄ · 7H ₂ O	0.02
	MnSO ₄ · 5H ₂ O	0.02
	Thiamine HCl	0.0002
	Yeast extract	1.0
25	CaCO ₃	30.0 (dry-heat-sterilized at 180°C for 2 h)

(Glucose and K₂HPO₄ separately sterilized)

The results are shown in Table 2. As shown in Table 2, the strains AJ13199/pYAHN, AJ13199/pYEAS, and AJ13199/pYFIK accumulated glutamic acid in a larger amount than the strain AJ13199/pUC21 in which an expression amount of amino acid excretion proteins was not enhanced.

Table 2.

Strain	Glutamic acid, g/l
AJ13199/pUC21	21.9
AJ13199/pYAHN	27.9
AJ13199/pYEAS	29.7
AJ13199/pYFIK	28.4

Example 4. Effect of *yeaS*, *yahN*, and *yfiK* DNA fragments amplification on lysine production.

(1). As the lysine-producing bacterium belonging to the genus *Escherichia*, *E. coli* strain W3110 (TyrA) described in European Patent Publication No. 488424 to which plasmid pCABD2 was introduced, described in International Publication No. WO 95/16042) was used. Specifically, plasmid pCABD2, and each of the plasmid pMW118::*yahN*, pMW118::*yeaS*, pMW118::*yfiK* and pMW118 were introduced to *E. coli* strain W3110 (TyrA) to obtain the following strains:

W3110 (tyrA)/pCABD2+pMW118::*yahN*

W3110 (tyrA)/pCABD2+pMW118::*yeaS*

W3110 (tyrA)/pCABD2+pMW118::*yfiK*

W3110 (tyrA)/pCABD2+pMW118.

Lysine productivity of these strains was estimated by culture. The composition of the used medium was as follows (g/l):

	Glucose	40.0
5	MgSO ₄ ·7H ₂ O	1.0
	(NH ₄) ₂ SO ₄	16.0
	K ₂ HPO ₄	1.0
	FeSO ₄ ·7H ₂ O	0.01
	MnSO ₄ ·7H ₂ O	0.01
10	Yeast extract (Difco)	2.0
	Tyrosine	0.1

Adjusted to pH 7.0 and autoclaved at 115° C for 10 minutes.

(Glucose and MgSO₄·7H₂O separately sterilized)

Pharmacopeial CaCO₃ 25 g/l (dry-heat-sterilized at
15 180°C for 2 h)

As antibiotics, 20 mg/l of streptomycin and 50 mg/l of ampicillin were added depending on a kind of a plasmid. Cultivation was conducted at 37°C for 30 hours with agitation at 115 rpm. The results are shown in Table 3.

20

Table 3

Strain	Lysine, g/l	Yield, (%)
W3110(<i>tyrA</i>)	0.08	0.2
W3110(<i>tyrA</i>)/pCABD2 + pMW118	12.2	30.5
W3110(<i>tyrA</i>)/pCABD2 + pMW118:: <i>yahN</i>	13.8	34.5
W3110(<i>tyrA</i>)/pCABD2 + pMW118:: <i>yeaS</i>	12.7	31.8
W3110(<i>tyrA</i>)/pCABD2 + pMW118:: <i>yfiK</i>	12.2	30.5

The result in Table 3 shows that the produced amount and the yield based on consumed sugar of lysine is increased by enhancement of YahN and YeaS.

(2). As the lysine-producing bacterium belonging to the genus *Escherichia*, *E. coli* strain VL614 was used. This strain is a derivative of the known *E. coli* strain VL613 (SU Patent No. 1354458). In turn, the strain VL613 was obtained from the known strain Gif102 (Theze, J. and Saint Girons. J.Bacteriol., 118, 990-998, 1974) in the three steps:

At the first step the mutants resistant to 2 mg/ml S-(2-aminoethyl)-L-cysteine were selected and among them the strain VL611 was found capable to produce L-lysine.

At the second step the genes involved in sucrose utilization and located on the transposon Tn2555 (Doroshenko et al., Mol. Biologiya, 22, 645-658, 1988), were introduced into VL611 using phage P1-mediated transduction giving the strain VL612.

At the third step, the mutation *rhtA*23 from the strain VKPM B-3996, conferring resistance to threonine and homoserine (US Patent No. 5,175,107) was introduced into VL612 by phage P1 transduction giving the strain VL613.

The *E. coli* strain VL614 was obtained by transduction of the wild-type allele of the *rhtA* gene from the *E. coli* strain VKPM B-6204 (MG1655 *zbi3058::Tn10*) to VL613. Transductants were selected on L-medium containing 10 mg/l tetracyclin, and among them the strain VL614 (*rhtA*⁺) sensitive to 10 g/l homoserine was found.

The strain VL614 was transformed with the pYGGA plasmid or with the pOK12 vector to obtain strains VL614/pYGGA (VKPM B-7719) and VL614/pOK12 (VKPM B-7722).

These strains were each cultivated at 37°C for 18 hours in a nutrient broth with 50 mg/l kanamycin, and 0.3 ml of the obtained culture was inoculated into 3 ml of a fermentation medium (Example 3) containing 0.3 g/l threonine, 0.3 g/l methionine and 50 mg/l kanamycin, in a 20 x 200 mm test tube, and cultivated at 37°C for 48 hours with a rotary shaker. After the cultivation, each accumulated amount of lysine and glutamate in the medium was determined by the known method.

The results are shown in Table 4.

Table 4

Strain	Lysine, g/l	Glutamate, g/l
VL614/pOK12	2.6	0.8
VL614/pYGGA	3.6	2.2

As shown in Table 4, the strain VL614/pYGGA accumulated lysine in a larger amount than the strain VL614/pOK12 in which the *yggA* gene was not enhanced. Besides, the strain VL614/pYGGA accumulated more glutamic acid than the strain VL614/pOK12.

Example 5. Effect of *yeaS*, *yahN*, and *yfiK* DNA fragments amplification on threonine, alanine, valine and isoleucine production.

As the threonine-producing bacterium belonging to the
5 genus *Escherichia*, the *E. coli* strain VL2054 was used. This strain was derived from the known *E. coli* strain VKPM B-3996 (US Patent No. 5,175,107) as follows.

Initially, a new recipient strain was constructed in several steps:

- 10 • The plasmidless derivative of the strain VKPM B-3996 was selected after spontaneous elimination of pVIC40 plasmid.
- The wild-type allele of the *rhtA* gene from the *E. coli* strain VKPM B-6204 (MG1655 *zbi3058::Tn10*) was introduced into the thus obtained strain by phage P1 mediated
15 transduction as in the Example 4.
- A mutation inactivating *kan* gene of the Tn5 transposon inserted into the *tdh* gene was obtained after NG mutagenesis and selection of kanamycin-sensitive cells still unable to degrade threonine. Thus the strain
20 VL2053 was obtained.

On the other hand, the threonine operon from pVIC40 was cloned into integrative Mud vector under the P_R promoter of the phage lambda. In addition, the *cat* gene of Tn9 conferring the resistance to chloramphenicol was cloned into
25 the same vector. The construct thus obtained was inserted into the chromosome of the *E. coli* strain C600 by use of the known method (US Patent No. 5,595,889) and transduced from

the thus obtained strain to VL2053, giving the new plasmidless threonine-producing strain VL2054. This strain accumulated in culture broth also alanine, valine and isoleucine.

5 The strain VL2054 was transformed with each of the plasmids pYEAS, pYFIK, and with the vector pUC21 to obtain *E. coli* strains VL2054/pYEAS (VKPM B-7707), VL2054/pYFIK (VKPM B-7712) and VL2054/pUC21 (VKPM B-7708).

10 These strains were each cultivated at 37°C for 18 hours in a nutrient broth with 100 mg/l ampicillin, and 0.3 ml of the obtained culture was inoculated into 3 ml of a fermentation medium (Example 3) containing 100 mg/l ampicillin, in a 20 x 200 mm test tube, and cultivated at 37°C for 48 hours with a rotary shaker. After the cultivation,
15 each accumulated amount of threonine, alanine, valine and isoleucine in the medium was determined by known method.

The results are shown in Table 5.

20 As shown in Table 5, the strain VL2054/pYFIK accumulated threonine in a larger amount than the strain VL2054/pUC21 in which the *yfiK* gene was not enhanced. Besides, the strain VL2054/pYEAS accumulated more alanine, valine and isoleucine than the strain VL2054/pUC21 in which the *yeas* gene was not enhanced.

Table 5

Strain	Amino acid accumulation, g/l			
	Threonine	Alanine	Valine	Isoleucine
VL2054/pUC21	5.8	0.4	0.31	0.15
VL2054/pYEAS	5.2	1.4	0.52	0.45
VL2054/pYFIK	8.8	0.5	0.22	0.14

Example 6. Effect of *yeaS* and *yfiK* DNA fragments amplification on histidine production.

5 As the histidine-producing bacterium belonging to the genus *Escherichia*, the strain *E. coli* VL2160 was used. This strain was obtained on the basis of the known strain NK5526 *hisG::Tn10* (VKPM B-3384) by phage P1-mediated transduction of the *hisG^R* mutation desensitizing ATP-phosphoribosyltransferase from the strain CC46
10 (Astvatsaturianz et al., Genetika, 24, 1928-1934, 1988). The strain *E. coli* VL2160 was transformed with each of the plasmids pYEAS, pYFIK, and with the vectors pUC21 to obtain *E. coli* strains VL2160/pYEAS (VKPM B-7753), *E. coli* VL2160/pYFIK (VKPM B-7754), *E. coli* VL2160/pUC21 (VKPM
15 B-7752).

These strains were each cultivated at 37°C for 18 hours in a nutrient broth with 100 mg/l ampicillin, and 0.3 ml of the obtained culture was inoculated into 3 ml of the
20 fermentation medium (Example 3) containing an increased amount of yeast extract (3 g/l) and 100 mg/l ampicillin, in a 20 x 200 mm test tube, and cultivated at 34°C for 68 hours with a rotary shaker.

After the cultivation, an accumulated amount of histidine in the medium was determined by known method. The results are shown in Table 6.

Table 6

Strain	Histidine, g/l
VL2160/pUC21	1.2
VL2160/pYEAS	1.8
VL2160/pYFIK	1.4

As shown in Table 6, the strains *E. coli* VL2160/ pYEAS and *E. coli* VL2160/pYFIK accumulated histidine in a larger amount than the strain *E. coli* VL2160/pUC21 in which the *yeaS* and *yfiK* genes were not enhanced.

Example 7. Effect of *yahN*, *yfiK* and *yeaS* DNA fragments amplification on proline production.

As the proline-producing bacterium belonging to the genus *Escherichia*, the strain VL2151 (W3350 *proB** Δ *putAP* Tn10) was used. This strain was obtained by transduction into W3350 of Δ *putAP* mutation linked to Tn10 and selecting tetracycline-resistant transductants unable to utilize proline as a sole carbon source. The thus obtained strain W3350 Δ *putAP* Tn10 was mutagenized with NG and mutants resistant to 20 mg/l of 3,4-dehydro-DL-proline were selected. Among them the strain VL2151 (W3350 *proB** Δ *putAP* Tn10) was found capable to produce proline.

The strain *E. coli* VL2151 was transformed with each of the plasmids pYEAS, pYFIK, pYAHN and with the vectors pUC21 to obtain *E. coli* strains VL2151/pYEAS (VKPM B-7714), VL2151/pYFIK (VKPM B-7713), VL2151/pYAHN (VKPM B-7748) and

5 *E. coli* VL2151/pUC21 (VKPM B-7715).

These strains were each cultivated at 37°C for 18 hours in a nutrient broth with 100 mg/l ampicillin, and 0.3 ml of the obtained culture was inoculated into 3 ml of a fermentation medium (Example 3) containing 100 mg/l

10 ampicillin, in a 20 x 200 mm test tube, and cultivated at 37°C for 48 hours with a rotary shaker. After the cultivation, an accumulated amount of proline in the medium was determined by known method. The results are shown in Table 7.

15 Table 7

Strain	Proline, g/l
VL2151/pUC21	1.8
VL2151/pYAHN	2.2
VL2151/pYEAS	2.1
VL2151/pYFIK	2.5

As shown in Table 7, the strains *E. coli* VL2151/pYFIK, *E. coli* VL2151/pYAHN and *E. coli* VL2151/pYEAS accumulated proline in a larger amount than the strain *E. coli* VL2151/pUC21 in which the *yfiK*, *yahN* and *yeaS* genes were not

20 enhanced. The amplification of *yfiK* gene had the most pronounced effect.

Example 8. Effect of *yggA* DNA fragments amplification on arginine production.

As arginine-producing bacterium belonging to the genus *Escherichia*, the strain W3350 *argE*::Tn10/pKA10 was used. This strain harbors a plasmid, pKA10, containing DNA region from *Corynebacterium* (*Brevibacterium*) *flavum* which complements at least *argA* and *argE* mutations in the recipient strain of *E. coli* K-12 (Kharitonov A. and Tarasov A.P. Molecular Genetics, Microbiology and Virology. No.9, 29-33, 1986).

The strain *E. coli* W3350 *argE*::Tn10/pKA10 was transformed with the plasmid pYGGA, or with the vector pOK12 to obtain the strains *E. coli* W3350 *argE*::Tn10/pKA10, pYGGA (VKPM B-7716) and *E. coli* W3350 *argE*::Tn10/pKA10, pOK12 (VKPM B-7718).

The thus obtained transformants were each cultivated at 37°C for 18 hours in a nutrient broth with 100 mg/l ampicillin and 50 mg/l kanamycin, and 0.3 ml of the obtained culture was inoculated into 3 ml of a fermentation medium (Example 3) containing 100 mg/l ampicillin and 50 mg/l kanamycin, in a 20 x 200 mm test tube, and cultivated at 37°C for 48 hours with a rotary shaker. After the cultivation, an accumulated amount of arginine in the medium was determined by known method.

The results are shown in Table 8.

Table 8

Strain	Arginine, g/l
W3350 <i>argE</i> ::Tn10/pKA10, pOK12	0.11
W3350 <i>argE</i> ::Tn10/pKA10, pYGGA	0.46

As shown in Table 8, the strains *E. coli* W3350
argE::Tn10/pKA10, pYGGA accumulated arginine in a larger
5 amount than the strain *E. coli* W3350 *argE*::Tn10/pKA10, pUC21
in which the *yggA* gene was not enhanced.

The following *E. coli* strains have been deposited
(according to international deposition based on Budapest
10 Treaty) in the Russian National Collection of Industrial
Microorganisms (VKPM) on December 29, 1998 under the
accession numbers shown in parenthesis.

AJ13199/pUC21 (VKPM B-7728)
AJ13199/pYAHN (VKPM B-7729)
15 AJ13199/pYEAS (VKPM B-7731)
AJ13199/pYFIK (VKPM B-7730)
VL614/pYGGA (VKPM B-7719)
VL614/pOK12 (VKPM B-7722)
VL2054/pYEAS (VKPM B-7707)
20 VL2054/pYFIK (VKPM B-7712)
VL2054/pUC21 (VKPM B-7708)
VL2160/pYEAS (VKPM B-7753)
VL2160/pYFIK (VKPM B-7754)
VL2160/pUC21 (VKPM B-7752)

VL2151/pUC21 (VKPM B-7715)

W3350 *argE*::Tn10/pKA10, pOK12 (VKPM B-7718)

SEQUENCE LISTING

<110> Ajinomoto Co., Inc.

<120> Method for Producing L-Amino Acid

<130>

<160> 24

<210> 1

<211> 27

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer for amplifying Escherichia coli yahN gene

<400> 1

ggcgagctcc cagtaaccgg aaataag

27

<210> 2

<211> 27

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer for amplifying Escherichia coli yahN gene

<400> 2

cgctctagaa aggaccacgc attacgg

27

<210> 3

<211> 27

<212> DNA

<223> Description of Artificial Sequence: primer for amplifying Escherichia coli yfiK gene

<400> 6
ggctctagat agcaagttac taagcgg 27

<210> 7
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: primer for amplifying Escherichia coli yggA gene

<400> 7
ctctgaattc tctcttatta gtttttctga ttgcc 35

<210> 8
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: primer for amplifying Escherichia coli yggA gene

<400> 8
cgtgacctgc agcgttctca cagcgcggta gcctttaa 38

<210> 9
<211> 672
<212> DNA
<213> Escherichia coli

<220>
<221> CDS
<222> (1)..(672)

<400> 9
atg atg cag tta gtt cac tta ttt atg gat gaa atc act atg gat cct 48
Met Met Gln Leu Val His Leu Phe Met Asp Glu Ile Thr Met Asp Pro
1 5 10 15
ttg cat gcc gtt tac ctg acc gta gga ctg ttc gtg att act ttt ttt 96
Leu His Ala Val Tyr Leu Thr Val Gly Leu Phe Val Ile Thr Phe Phe

```
<210> 10
<211> 223
<212> PRT
<213> Escherichia coli
```

<400> 10

```

Met Met Gln Leu Val His Leu Phe Met Asp Glu Ile Thr Met Asp Pro
 1           5           10           15
Leu His Ala Val Tyr Leu Thr Val Gly Leu Phe Val Ile Thr Phe Phe
          20           25           30
Asn Pro Gly Ala Asn Leu Phe Val Val Val Gln Thr Ser Leu Ala Ser
          35           40           45
Gly Arg Arg Ala Gly Val Leu Thr Gly Leu Gly Val Ala Leu Gly Asp
          50           55           60
Ala Phe Tyr Ser Gly Leu Gly Leu Phe Gly Leu Ala Thr Leu Ile Thr
          65           70           75           80
Gln Cys Glu Glu Ile Phe Ser Leu Ile Arg Ile Val Gly Gly Ala Tyr
          85           90           95
Leu Leu Trp Phe Ala Trp Cys Ser Met Arg Arg Gln Ser Thr Pro Gln
          100          105          110
Met Ser Thr Leu Gln Gln Pro Ile Ser Ala Pro Trp Tyr Val Phe Phe
          115          120          125
Arg Arg Gly Leu Ile Thr Asp Leu Ser Asn Pro Gln Thr Val Leu Phe
          130          135          140
Phe Ile Ser Ile Phe Ser Val Thr Leu Asn Ala Glu Thr Pro Thr Trp
          145          150          155          160
Ala Arg Leu Met Ala Trp Ala Gly Ile Val Leu Ala Ser Ile Ile Trp
          165          170          175
Arg Val Phe Leu Ser Gln Ala Phe Ser Leu Pro Ala Val Arg Arg Ala
          180          185          190
Tyr Gly Arg Met Gln Arg Val Ala Ser Arg Val Ile Gly Ala Ile Ile
          195          200          205
Gly Val Phe Ala Leu Arg Leu Ile Tyr Glu Gly Val Thr Gln Arg
          210          215          220

```

<210> 11

<211> 639

<212> DNA

<213> Escherichia coli

<220>

<221> CDS

<222> (1)..(639)

<400> 11

gtg ttc gct gaa tac ggg gtt ctg aat tac tgg acc tat ctg gtt ggg 48

Met	Phe	Ala	Glu	Tyr	Gly	Val	Leu	Asn	Tyr	Trp	Thr	Tyr	Leu	Val	Gly	
1				5					10					15		
gcc	att	ttt	att	gtg	ttg	gtg	cca	ggg	cca	aat	acc	ctg	ttt	gta	ctc	96
Ala	Ile	Phe	Ile	Val	Leu	Val	Pro	Gly	Pro	Asn	Thr	Leu	Phe	Val	Leu	
			20					25					30			
aaa	aat	agc	gtc	agt	agc	ggg	atg	aaa	ggc	ggg	tat	ctt	gcg	gcc	tgc	144
Lys	Asn	Ser	Val	Ser	Ser	Gly	Met	Lys	Gly	Gly	Tyr	Leu	Ala	Ala	Cys	
		35					40				45					
ggg	gta	ttt	att	ggc	gat	gcg	gta	ttg	atg	ttt	ctg	gca	tgg	gct	gga	192
Gly	Val	Phe	Ile	Gly	Asp	Ala	Val	Leu	Met	Phe	Leu	Ala	Trp	Ala	Gly	
	50					55					60					
gtg	gcg	aca	tta	att	aag	acc	acc	ccg	ata	tta	ttc	aac	att	gta	cgt	240
Val	Ala	Thr	Leu	Ile	Lys	Thr	Thr	Pro	Ile	Leu	Phe	Asn	Ile	Val	Arg	
	65				70				75				80			
tat	ctt	ggg	gcg	ttt	tat	ttg	ctc	tat	ctg	ggg	agt	aaa	att	ctt	tac	288
Tyr	Leu	Gly	Ala	Phe	Tyr	Leu	Leu	Tyr	Leu	Gly	Ser	Lys	Ile	Leu	Tyr	
			85					90				95				
gcg	acc	ctg	aag	ggg	aaa	aat	agc	gag	gcc	aaa	tcc	gat	gag	ccc	caa	336
Ala	Thr	Leu	Lys	Gly	Lys	Asn	Ser	Glu	Ala	Lys	Ser	Asp	Glu	Pro	Gln	
		100						105				110				
tac	ggg	gct	att	ttt	aaa	cgc	gcg	tta	att	ttg	agc	ctg	act	aat	ccg	384
Tyr	Gly	Ala	Ile	Phe	Lys	Arg	Ala	Leu	Ile	Leu	Ser	Leu	Thr	Asn	Pro	
	115					120					125					
aaa	gcc	att	ttg	ttc	tat	gtg	tcg	ttt	ttc	gta	cag	ttt	atc	gat	gtt	432
Lys	Ala	Ile	Leu	Phe	Tyr	Val	Ser	Phe	Phe	Val	Gln	Phe	Ile	Asp	Val	
	130					135				140						
aat	gcc	cca	cat	acg	gga	att	tca	ttc	ttt	att	ctg	gcg	gcg	acg	ctg	480
Asn	Ala	Pro	His	Thr	Gly	Ile	Ser	Phe	Phe	Ile	Leu	Ala	Ala	Thr	Leu	
	145				150				155				160			
gaa	ctg	gtg	agt	ttc	tgc	tat	ttg	agc	ttc	ctg	att	ata	tct	ggg	gct	528
Glu	Leu	Val	Ser	Phe	Cys	Tyr	Leu	Ser	Phe	Leu	Ile	Ile	Ser	Gly	Ala	
			165					170				175				
ttt	gtc	acg	cag	tac	ata	cgt	acc	aaa	aag	aaa	ctg	gct	aaa	gtt	ggc	576
Phe	Val	Thr	Gln	Tyr	Ile	Arg	Thr	Lys	Lys	Lys	Leu	Ala	Lys	Val	Gly	
		180						185				190				
aac	tca	ctg	att	ggg	ttg	atg	ttc	gtg	ggg	ttc	gct	gcc	cga	ctg	gcg	624
Asn	Ser	Leu	Ile	Gly	Leu	Met	Phe	Val	Gly	Phe	Ala	Ala	Arg	Leu	Ala	
		195					200					205				
acg	ctg	caa	tcc	tga												639
Thr	Leu	Gln	Ser													
	210															

<210> 12
 <211> 212
 <212> PRT
 <213> Escherichia coli

<400> 12

```

Met Phe Ala Glu Tyr Gly Val Leu Asn Tyr Trp Thr Tyr Leu Val Gly
 1           5           10           15
Ala Ile Phe Ile Val Leu Val Pro Gly Pro Asn Thr Leu Phe Val Leu
           20           25           30
Lys Asn Ser Val Ser Ser Gly Met Lys Gly Gly Tyr Leu Ala Ala Cys
           35           40           45
Gly Val Phe Ile Gly Asp Ala Val Leu Met Phe Leu Ala Trp Ala Gly
           50           55           60
Val Ala Thr Leu Ile Lys Thr Thr Pro Ile Leu Phe Asn Ile Val Arg
           65           70           75           80
Tyr Leu Gly Ala Phe Tyr Leu Leu Tyr Leu Gly Ser Lys Ile Leu Tyr
           85           90           95
Ala Thr Leu Lys Gly Lys Asn Ser Glu Ala Lys Ser Asp Glu Pro Gln
           100          105          110
Tyr Gly Ala Ile Phe Lys Arg Ala Leu Ile Leu Ser Leu Thr Asn Pro
           115          120          125
Lys Ala Ile Leu Phe Tyr Val Ser Phe Phe Val Gln Phe Ile Asp Val
           130          135          140
Asn Ala Pro His Thr Gly Ile Ser Phe Phe Ile Leu Ala Ala Thr Leu
           145          150          155          160
Glu Leu Val Ser Phe Cys Tyr Leu Ser Phe Leu Ile Ile Ser Gly Ala
           165          170          175
Phe Val Thr Gln Tyr Ile Arg Thr Lys Lys Lys Leu Ala Lys Val Gly
           180          185          190
Asn Ser Leu Ile Gly Leu Met Phe Val Gly Phe Ala Ala Arg Leu Ala
           195          200          205
Thr Leu Gln Ser
           210

```

<210> 13
 <211> 588
 <212> DNA
 <213> Escherichia coli

<220>
 <221> CDS

<400> 13

[illegible]

<400> 15
 gtg ttt tct tat tac ttt caa ggt ctt gca ctt ggg gcg gct atg atc 48
 Met Phe Ser Tyr Tyr Phe Gln Gly Leu Ala Leu Gly Ala Ala Met Ile
 1 5 10 15
 cta ccg ctc ggt cca caa aat gct ttt gtg atg aat cag ggc ata cgt 96
 Leu Pro Leu Gly Pro Gln Asn Ala Phe Val Met Asn Gln Gly Ile Arg
 20 25 30
 cgt cag tac cac att atg att gcc tta ctt tgt gct atc agc gat ttg 144
 Arg Gln Tyr His Ile Met Ile Ala Leu Leu Cys Ala Ile Ser Asp Leu
 35 40 45
 gtc ctg att tgc gcc ggg att ttt ggt ggc agc gcg tta ttg atg cag 192
 Val Leu Ile Cys Ala Gly Ile Phe Gly Gly Ser Ala Leu Leu Met Gln
 50 55 60
 tcg ccg tgg ttg ctg gcg ctg gtc acc tgg ggc ggc gta gcc ttc ttg 240
 Ser Pro Trp Leu Leu Ala Leu Val Thr Trp Gly Gly Val Ala Phe Leu
 65 70 75 80
 ctg tgg tat ggt ttt ggc gct ttt aaa aca gca atg agc agt aat att 288
 Leu Trp Tyr Gly Phe Gly Ala Phe Lys Thr Ala Met Ser Ser Asn Ile
 85 90 95
 gag tta gcc agc gcc gaa gtc atg aag caa ggc aga tgg aaa att atc 336
 Glu Leu Ala Ser Ala Glu Val Met Lys Gln Gly Arg Trp Lys Ile Ile
 100 105 110
 gcc acc atg ttg gca gtg acc tgg ctg aat ccg cat gtt tac ctg gat 384
 Ala Thr Met Leu Ala Val Thr Trp Leu Asn Pro His Val Tyr Leu Asp
 115 120 125
 act ttt gtt gta ctg ggc agc ctt ggc ggg caa ctt gat gtg gaa cca 432
 Thr Phe Val Val Leu Gly Ser Leu Gly Gly Gln Leu Asp Val Glu Pro
 130 135 140
 aaa cgc tgg ttt gca ctc ggg aca att agc gcc tct ttc ctg tgg ttc 480
 Lys Arg Trp Phe Ala Leu Gly Thr Ile Ser Ala Ser Phe Leu Trp Phe
 145 150 155 160
 ttt ggt ctg gct ctt ctc gca gcc tgg ctg gca ccg cgt ctg cgc acg 528
 Phe Gly Leu Ala Leu Leu Ala Ala Trp Leu Ala Pro Arg Leu Arg Thr
 165 170 175
 gca aaa gca cag cgc att atc aat ctg gtt gtg gga tgt gtt atg tgg 576
 Ala Lys Ala Gln Arg Ile Ile Asn Leu Val Val Gly Cys Val Met Trp
 180 185 190
 ttt att gcc ttg cag ctg gcg aga gac ggt att gct cat gca caa gcc 624
 Phe Ile Ala Leu Gln Leu Ala Arg Asp Gly Ile Ala His Ala Gln Ala
 195 200 205
 ttg ttc agt tag 636

Leu Phe Ser
210

<210> 16
<211> 211
<212> PRT
<213> Escherichia coli

<400> 16
Met Phe Ser Tyr Tyr Phe Gln Gly Leu Ala Leu Gly Ala Ala Met Ile
1 5 10 15
Leu Pro Leu Gly Pro Gln Asn Ala Phe Val Met Asn Gln Gly Ile Arg
20 25 30
Arg Gln Tyr His Ile Met Ile Ala Leu Leu Cys Ala Ile Ser Asp Leu
35 40 45
Val Leu Ile Cys Ala Gly Ile Phe Gly Gly Ser Ala Leu Leu Met Gln
50 55 60
Ser Pro Trp Leu Leu Ala Leu Val Thr Trp Gly Gly Val Ala Phe Leu
65 70 75 80
Leu Trp Tyr Gly Phe Gly Ala Phe Lys Thr Ala Met Ser Ser Asn Ile
85 90 95
Glu Leu Ala Ser Ala Glu Val Met Lys Gln Gly Arg Trp Lys Ile Ile
100 105 110
Ala Thr Met Leu Ala Val Thr Trp Leu Asn Pro His Val Tyr Leu Asp
115 120 125
Thr Phe Val Val Leu Gly Ser Leu Gly Gly Gln Leu Asp Val Glu Pro
130 135 140
Lys Arg Trp Phe Ala Leu Gly Thr Ile Ser Ala Ser Phe Leu Trp Phe
145 150 155 160
Phe Gly Leu Ala Leu Leu Ala Ala Trp Leu Ala Pro Arg Leu Arg Thr
165 170 175
Ala Lys Ala Gln Arg Ile Ile Asn Leu Val Val Gly Cys Val Met Trp
180 185 190
Phe Ile Ala Leu Gln Leu Ala Arg Asp Gly Ile Ala His Ala Gln Ala
195 200 205
Leu Phe Ser
210

<210> 17
<211> 20
<212> DNA
<213> Artificial Sequence

0460460

<223> Description of Artificial Sequence: primer for amplifying Escherichia coli yahN gene

gtgtggaacc gacgccgat 20

<213> Artificial Sequence

<223> Description of Artificial Sequence: primer for amplifying Escherichia coli yahN gene

tgttgtatgg tacggggttc gag 23

<213> Artificial Sequence

<223> Description of Artificial Sequence: primer for amplifying Escherichia coli yeaS gene

ctttgccaat cccgtctccc 20

<213> Artificial Sequence

<223> Description of Artificial Sequence: primer for amplifying Escherichia coli yeaS gene

<400> 20

20

<213> Artificial Sequence

<223> Description of Artificial Sequence: primer for amplifying Escherichia coli yfiK gene

26

<213> Artificial Sequence

<223> Description of Artificial Sequence: primer for amplifying Escherichia coli yfiK gene

20

<213> Artificial Sequence

<223> Description of Artificial Sequence: primer for amplifying Escherichia coli yggA gene

21

<213> Artificial Sequence

<223> Description of Artificial Sequence: primer for amplifying Escherichia coli yggA gene

ggcaagctta gcgcctctgt t

21

What is claimed is:

1. A bacterium belonging to the genus *Escherichia* and having an ability to produce an L-amino acid, wherein the ability to produce the L-amino acid is increased by
5 increasing an expression amount of at least one protein selected from the group consisting of the following proteins of (A) to (H):

(A) a protein having an amino acid sequence shown in SEQ ID NO: 10 in Sequence Listing;

10 (B) a protein which has an amino acid sequence including deletion, substitution, insertion, addition or inversion of one or several amino acids in the amino acid sequence shown in SEQ ID NO: 10 in Sequence Listing, and which has an activity of increasing the ability to produce the
15 L-amino acid of the bacterium having the protein;

(C) a protein having an amino acid sequence shown in SEQ ID NO: 12 in Sequence Listing;

(D) a protein which has an amino acid sequence including deletion, substitution, insertion, addition or
20 inversion of one or several amino acids in the amino acid sequence shown in SEQ ID NO: 12 in Sequence Listing, and which has an activity of increasing the ability to produce the L-amino acid of the bacterium having the protein;

(E) a protein having an amino acid sequence shown in
25 SEQ ID NO: 14 in Sequence Listing;

(F) a protein which has an amino acid sequence including deletion, substitution, insertion, addition or inversion of one or several amino acids in the amino acid sequence shown in SEQ ID NO: 14 in Sequence Listing, and which
30 has an activity of increasing the ability to produce the

L-amino acid of the bacterium having the protein;

(G) a protein having an amino acid sequence shown in SEQ ID NO: 16 in Sequence Listing; or

(H) a protein which has an amino acid sequence.

5 including deletion, substitution, insertion, addition or inversion of one or several amino acids in the amino acid sequence shown in SEQ ID NO: 16 in Sequence Listing, and which has an activity of increasing the ability to produce the L-amino acid of the bacterium having the protein.

10 2. The bacterium according to claim 1, wherein said L-amino acid is L-lysine and an expression amount of at least one protein selected from the group consisting of said proteins (A) to (D), (G) and (H) is increased.

15 3. The bacterium according to claim 1, wherein said L-amino acid is L-glutamic acid and an expression amount of at least one protein selected from the group consisting of said proteins (A) to (H) is increased.

20 4. The bacterium according to claim 1, wherein said L-amino acid is L-alanine and an expression amount of at least one protein selected from the group consisting of said proteins (C) and (D) is increased.

25 5. The bacterium according to claim 1, wherein said L-amino acid is L-valine and an expression amount of at least one protein selected from the group consisting of said proteins (C) and (D) is increased.

6. The bacterium according to claim 1, wherein said L-amino acid is L-histidine and an expression amount of at least one protein selected from the group consisting of said proteins (C) to (F) is increased.

30 7. The bacterium according to claim 1, wherein said L-

amino acid is L-proline and an expression amount of at least one protein selected from the group consisting of said proteins (A) to (F) is increased.

8. The bacterium according to claim 1, wherein said L-amino acid is L-threonine and an expression amount of at least one protein selected from the group consisting of said proteins (E) and (F) is increased.

9. The bacterium according to claim 1, wherein said L-amino acid is L-arginine and an expression amount of at least one protein selected from the group consisting of said proteins (G) and (H) is increased.

10. The bacterium according to claim 1, wherein said L-amino acid is L-isoleucine and an expression amount of at least one protein selected from the group consisting of said proteins (C) and (D) is increased.

11. The bacterium according to any one of claims 1 to 10, wherein a copy number of a DNA coding for said protein in a cell is increased.

12. The bacterium according to claim 11, wherein said DNA is carried on a multicopy vector in the cell.

13. The bacterium according to claim 11, wherein said DNA is carried on a transposon in the cell.

14. A method for producing an L-amino acid, comprising the steps of:

25 cultivating the bacterium as defined in claim 1 in a culture medium, to produce and accumulate the L-amino acid in the medium, and

 recovering the L-amino acid from the medium.

15. The method according to claim 14, wherein said L-amino acid is L-lysine and said bacterium is one in which an

expression amount of at least one protein selected from the group consisting of said proteins (A) to (D), (G) and (H) is increased.

5 16. The method according to claim 14, wherein said L-amino acid is L-glutamic acid and said bacterium is one in which an expression amount of at least one protein selected from the group consisting of said proteins (A) to (H) is increased.

10 17. The method according to claim 14, wherein said L-amino acid is L-alanine and said bacterium is one in which an expression amount of at least one protein selected from the group consisting of said proteins (C) and (D) is increased.

15 18. The method according to claim 14, wherein said L-amino acid is L-valine and said bacterium is one in which an expression amount of at least one protein selected from the group consisting of said proteins (C) and (D) is increased.

20 19. The method according to claim 14, wherein said L-amino acid is L-histidine and said bacterium is one in which an expression amount of at least one protein selected from the group consisting of said proteins (C) to (F) is increased.

20. The method according to claim 14, wherein said L-amino acid is L-proline and said bacterium is one in which an expression amount of at least one protein selected from the group consisting of said proteins (A) to (F) is increased.

25 21. The method according to claim 14, wherein said L-amino acid is L-threonine and said bacterium is one in which an expression amount of at least one protein selected from the group consisting of said proteins (E) and (F) is increased.

30 22. The method according to claim 14, wherein said L-amino acid is L-arginine and said bacterium is one in which an

23. The method according to claim 14, wherein said L-amino acid is L-isoleucine and said bacterium is one in which an

24. The method according to any one of claims 14 to 23, wherein a copy number of a DNA coding for said protein in a cell of said bacterium is increased.

26. The method according to claim 24, wherein said DNA is carried on a transposon in the cell.

Abstract of the Disclosure

A bacterium belonging to the genus *Escherichia* and having an ability to produce an L-amino acid, wherein the
5 ability to produce the L-amino acid is increased by increasing an expression amount of an L-amino acid excretion protein, and a method for producing the L-amino acid using the bacterium.

Declaration, Power Of Attorney and Petition

Page 1 of 3

WE (I) the undersigned inventor(s), hereby declare(s) that:

My residence, post office address and citizenship are as stated below next to my name,

We (I) believe that we are (I am) the original, first, and joint (sole) inventor(s) of the subject matter which is claimed and for which a patent is sought on the invention entitled

METHOD FOR PRODUCING L-AMINO ACID

the specification of which

☒ is attached hereto.

☐ was filed on _____ as

Application Serial No. _____

and amended on _____.

☐ was filed as PCT international application

Number _____

on _____,

and was amended under PCT Article 19

on _____ (if applicable).

We (I) hereby state that we (I) have reviewed and understand the contents of the above-identified specification, including the claims, as amended by any amendment referred to above.

We (I) acknowledge the duty to disclose information known to be material to the patentability of this application as defined in Section 1.56 of Title 37 Code of Federal Regulations.

We (I) hereby claim foreign priority benefits under 35 U.S.C. § 119(a)-(d) or § 365(b) of any foreign application(s) for patent or inventor's certificate, or § 365(a) of any PCT International application which designated at least one country other than the United States, listed below and have also identified below, by checking the box, any foreign application for patent or inventor's certificate, or PCT International application having a filing date before that of the application on which priority is claimed. Prior Foreign Application(s)

Application No.	Country	Day/Month/Year	Priority Claimed	
98124016	Russia	30/12/1998	<input checked="" type="checkbox"/> Yes	<input type="checkbox"/> No
99104431	Russia	09/03/1999	<input checked="" type="checkbox"/> Yes	<input type="checkbox"/> No
_____	_____	_____	<input type="checkbox"/> Yes	<input type="checkbox"/> No
_____	_____	_____	<input type="checkbox"/> Yes	<input type="checkbox"/> No

We (I) hereby claim the benefit under Title 35, United States Code, § 119(e) of any United States provisional application(s) listed below.

(Application Number)

(Filing Date)

(Application Number)

(Filing Date)

We (I) hereby claim the benefit under 35 U.S.C. § 120 of any United States application(s), or § 365(c) of any PCT International application designating the United States, listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States or PCT International application in the manner provided by the first paragraph of 35 U.S.C. § 112, I acknowledge the duty to disclose information which is material to patentability as defined in 37 CFR § 1.56 which became available between the filing date of the prior application and the national or PCT International filing date of this application.

Application Serial No.

Filing Date

Status (pending, patented,
abandoned)

Application Serial No.	Filing Date	Status (pending, patented, abandoned)


And we (I) hereby appoint: Norman F. Oblon, Registration Number 24,618; Marvin J. Spivak, Registration Number 24,913; C. Irvin McClelland, Registration Number 21,124; Gregory J. Maier, Registration Number 25,599; Arthur I. Neustadt, Registration Number 24,854; Richard D. Kelly, Registration Number 27,757; James D. Hamilton, Registration Number 28,421; Eckhard H. Kuesters, Registration Number 28,870; Robert T. Pous, Registration Number 29,099; Charles L. Gholz, Registration Number 26,395; Vincent J. Sunderdick, Registration Number 29,004; William E. Beaumont, Registration Number 30,996; Steven B. Kelber, Registration Number 30,073; Robert F. Gnuse, Registration Number 27,295; Jean-Paul Lavalleye, Registration Number 31,451; Timothy R. Schwartz, Registration Number 32,171; Stephen G. Baxter, Registration Number 32,884; Martin M. Zoltick, Registration Number 35,745; Robert W. Hahl, Registration Number 33,893; Richard L. Treanor, Registration Number 36,379; Steven P. Weihrouch, Registration Number 32,829; John T. Goolkasian, Registration Number 26,142; Marc R. Labgold, Registration Number 34,651; William J. Healey, Registration Number 36,160; and Richard L. Chinn, Registration Number 34,305; our (my) attorneys, with full powers of substitution and revocation, to prosecute this application and to transact all business in the Patent Office connected therewith; and we (I) hereby request that all correspondence regarding this application be sent to the firm of OBLON, SPIVAK, MCCLELLAND, MAIER & NEUSTADT, P.C., whose Post Office Address is: Fourth Floor, 1755 Jefferson Davis Highway, Arlington, Virginia 22202.

We (I) declare that all statements made herein of our (my) own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issuing thereon.

Vitaliy Arkadievich LIVSHITS
NAME OF FIRST SOLE INVENTOR

Residence: Moscow Russia

Citizen: Russia


Signature of Inventor

Post Office Address: kv. 84, korpus 1,
Sumskoy proezd 5, Moscow, 113208,

November 24, 1999
Date

Natalia Pavlovna ZAKATAEVA

NAME OF SECOND JOINT INVENTOR

Residence: Moscow Russia

Zakataeva

Signature of Inventor

Citizen: Russia

Post Office Address: kv. 27, korpus 8, ulitsa
Novatorov 40, Moscow, 117421, Russia

November 24, 1999

Date

Kazuo NAKANISHI

NAME OF THIRD JOINT INVENTOR

Residence: Yokohama-shi, Kanagawa, Japan

Nakanishi

Signature of Inventor

Citizen: Japanese

Post Office Address: Kuden-Haitsu 1-126,
Kuden-Cho 873-5, Sakae-ku, Yokohama-shi
247-0014 Japan,

November 24, 1999

Date

Vladimir Veniaminovich ALESHIN

NAME OF FOURTH JOINT INVENTOR

Redsidence: Moscow Russia

Aleshin

Signature of Inventor

Citizen of: Russia

Post Office Address: pos Institute 6,
Kaluga region, Borovsk, 249010, Russia

November 24, 1999

Date

Petr Vladimirovich TROSHIN

NAME OF FIFTH JOINT INVENTOR

Residence: Moscow Russia

Troshin

Signature of Inventor

Citizen of: Russia

Post Office Address: kv. 129, Shipilovskiy
pr. 67, Moscow, 115561, Russia

November 24, 1999

Date

Irina Lyvovna TOKHMAKOVA

NAME OF SIXTH JOINT INVENTOR

Residence: Moscow Russia

Citizen of: Russia

Irina
Signature of Inventor

Post Office Address: kv. 26, s/p Vysotka 1,
Isrtrinskiy r-n, Moscow region, 143513, Russia

November 24, 1999

Date

NAME OF SEVENTH JOINT INVENTOR

Residence: _____

Signature of Inventor

Citizen of: _____

Post Office Address: _____

Date

NAME OF EIGHTH JOINT INVENTOR

Residence: _____

Signature of Inventor

Citizen of: _____

Post Office Address: _____

Date

NAME OF NINTH JOINT INVENTOR

Residence: _____

Signature of Inventor

Citizen of: _____

Post Office Address: _____

Date

**Budapest treaty on the international
recognition of the deposit of microorganisms
for the purposes of patent procedure**

To
State Scientific Centre of
Russian Federation
GNIIGENETIKA
Moscow 113545
1-st Dorozhny proezd 1 Russia
name and address
of depositor

INTERNATIONAL FORM
receipt in the case of an original deposit
issued pursuant to Rule 7.1 by the
INTERNATIONAL DEPOSITARY AUTHORITY
identified at the bottom of this page

I. IDENTIFICATION OF THE MICROORGANISM

Identification reference given by the
DEPOSITOR

Escherichia coli VL2054(pYEAS)

Accession number given by the
INTERNATIONAL DEPOSITARY-
AUTHORITY:
VKPM B- 7707

II. SCIENTIFIC DESCRIPTION AND/OR PROPOSED TAXONOMIC DESIGNATION

The microorganism identified under I above was accompanied by:

☒ a scientific description

Producer of threonine

☐ a proposed taxonomic designation
(Mark with a cross where applicable)

III. RECEIPT AND ACCEPTANCE

This International Depositary Authority accepts the microorganism identified under I above,
1 which was received by it on 29.12.1998 (date of original deposit)

IV. RECEIPT OF REQUEST FOR CONVERSION

The microorganism identified under I above was received by this International Depositary
Authority
on (date of original deposit) and a request to convert
the
original deposit to a deposit under the Budapest Treaty was received by it on
(date of receipt of request for conversion)

V. INTERNATIONAL DEPOSITARY AUTHORITY


Name: Russian National Collection
of Industrial Microorganisms (VKPM)
Depositary
GNIIGenetika
Address: Russia 113545 Moscow
1 Dorozhny proezd 1

Signature (s) of person(s) having the
power to represent the International

Authority or authorized official(s):

Date: Sineoky S.P.

1 Where Rule 6.4(d) applies, such date is the date on which the status of international
depositary authority was acquired



Appendix 3
page 14

Budapest treaty on the international
recognition of the deposit of microorganisms
for the purposes of patent procedure

To
State Scientific Centre of
Russian Federation
GNIIGENETIKA
Moscow 113545
1-st Dorozhny proezd 1 Russia
name and address
of depositor

INTERNATIONAL FORM
receipt in the case of an original deposit
issued pursuant to Rule 7.1 by the
INTERNATIONAL DEPOSITARY AUTHORITY
identified at the bottom of this page

I. IDENTIFICATION OF THE MICROORGANISM

Identification reference given by the DEPOSITOR	Accession number given by the INTERNATIONAL DEPOSITARY- AUTHORITY:
Escherichia coli VL2054 (pUC21)	VKPM B-7708

II. SCIENTIFIC DESCRIPTION AND/OR PROPOSED TAXONOMIC DESIGNATION

The microorganism identified under I above was accompanied by:

☒ a scientific description Producer of threonine, alanine
☐ a proposed taxonomic designation
(Mark with a cross where applicable)

III. RECEIPT AND ACCEPTANCE

This International Depositary Authority accepts the microorganism identified under I above,
which was received by it on 29.12.1998 (date of original deposit)

IV. RECEIPT OF REQUEST FOR CONVERSION

The microorganism identified under I above was received by this International Depositary
Authority on (date of original deposit) and a request to convert
the original deposit to a deposit under the Budapest Treaty was received by it on
(date of receipt of request for conversion)


V. INTERNATIONAL DEPOSITARY AUTHORITY

Name: Russian National Collection
of Industrial Microorganisms (VKPM)
Depositary
GNIIGenetika
Address: Russia 113545 Moscow
1 Dorozhny proezd 1

Signature (s) of person(s) having the
power to represent the International

Authority or authorized official(s):
Sinecky S.P.

1 Where Rule 6.4(d) applies, such date is the date on which the status of international
depository authority was acquired



Appendix 3
page 14

Budapest treaty on the international
recognition of the deposit of microorganisms
for the purposes of patent procedure

To
State Scientific Centre of
Russian Federation
GNIIGENETIKA
Moscow 113545
1-st Dorozhny proezd 1 Russia
name and address
of depositor

INTERNATIONAL FORM
receipt in the case of an original deposit
issued pursuant to Rule 7.1 by the
INTERNATIONAL DEPOSITARY AUTHORITY
identified at the bottom of this page

I. IDENTIFICATION OF THE MICROORGANISM

Identification reference given by the
DEPOSITOR

Escherichia coli VL2054(pYFIK)

Accession number given by the
INTERNATIONAL DEPOSITARY-
AUTHORITY:
VKPM B- 7712

II. SCIENTIFIC DESCRIPTION AND/OR PROPOSED TAXONOMIC DESIGNATION

The microorganism identified under I above was accompanied by:

☒ a scientific description Producer of threonine

☐ a proposed taxonomic designation
(Mark with a cross where applicable)

III. RECEIPT AND ACCEPTANCE

This International Depositary Authority accepts the microorganism identified under I above,
I which was received by it on 29.12.1998 (date of original deposit)

IV. RECEIPT OF REQUEST FOR CONVERSION

The microorganism identified under I above was received by this International Depositary
Authority on (date of original deposit) and a request to convert
the original deposit to a deposit under the Budapest Treaty was received by it on
(date of receipt of request for conversion)


V. INTERNATIONAL DEPOSITARY AUTHORITY

Name: Russian National Collection
of Industrial Microorganisms (VKPM)
Depositary
GNIIGenetika
Address: Russia 113545 Moscow
1 Dorozhny proezd 1

Signature (s) of person(s) having the
power to represent the International

Authority or authorized official(s):
Date: Simeoky S.P.

I Where Rule 6.4(d) applies, such date is the date on which the status of international
depositary authority was acquired



Appendix 3

page 14

Budapest treaty on the international
recognition of the deposit of microorganisms
for the purposes of patent procedure

To
State Scientific Centre of
Russian Federation
GNIIGENETIKA
Moscow 113545
1-st Dorozhny proezd 1 Russia
name and address
of depositor

INTERNATIONAL FORM
receipt in the case of an original deposit
issued pursuant to Rule 7.1 by the
INTERNATIONAL DEPOSITARY AUTHORITY
identified at the bottom of this page

I. IDENTIFICATION OF THE MICROORGANISM

Identification reference given by the DEPOSITOR	Accession number given by the INTERNATIONAL DEPOSITARY- AUTHORITY:
Escherichia coli VL2151 (pYFIK)	VKPM B- 7713

II. SCIENTIFIC DESCRIPTION AND/OR PROPOSED TAXONOMIC DESIGNATION

The microorganism identified under I above was accompanied by:

☒ a scientific description Producer of proline

☐ a proposed taxonomic designation
(Mark with a cross where applicable)

III. RECEIPT AND ACCEPTANCE

This International Depositary Authority accepts the microorganism identified under I above,
1 which was received by it on 29.12.1998 (date of original deposit)

IV. RECEIPT OF REQUEST FOR CONVERSION

The microorganism identified under I above was received by this International Depositary
Authority
on (date of original deposit) and a request to convert
the
original deposit to a deposit under the Budapest Treaty was received by it on
(date of receipt of request for conversion)


V. INTERNATIONAL DEPOSITARY AUTHORITY

Name: Russian National Collection
of Industrial Microorganisms (VKPM)
Depositary
GNIIGenetika
Address: Russia 113545 Moscow
1 Dorozhny proezd 1

Signature (s) of person(s) having the
power to represent the International

Authority or authorized official(s):
Date: Sineoky S.P.

1 Where Rule 6.4(d) applies, such date is the date on which the status of international
depositary authority was acquired



Handwritten signature of Sineoky S.P. and a circular stamp of the International Depositary Authority.

045573-12399

Appendix 3
page 14

Budapest treaty on the international
recognition of the deposit of microorganisms
for the purposes of patent procedure

To
State Scientific Centre of
Russian Federation
GNIIGENETIKA
Moscow 113545
1-st Dorozhny proezd 1 Russia
name and address
of depositor

INTERNATIONAL FORM
receipt in the case of an original deposit
issued pursuant to Rule 7.1 by the
INTERNATIONAL DEPOSITARY AUTHORITY
identified at the bottom of this page

I. IDENTIFICATION OF THE MICROORGANISM

Identification reference given by the DEPOSITOR	Accession number given by the INTERNATIONAL DEPOSITARY- AUTHORITY:
Escherichia coli VL2151 (pYEAS)	VKPM B- 7714

II. SCIENTIFIC DESCRIPTION AND/OR PROPOSED TAXONOMIC DESIGNATION

The microorganism identified under I above was accompanied by:

☒ a scientific description Producer of proline

☐ a proposed taxonomic designation
(Mark with a cross where applicable)

III. RECEIPT AND ACCEPTANCE

This International Depositary Authority accepts the microorganism identified under I above,
I which was received by it on 29.12.1998 (date of original deposit)

IV. RECEIPT OF REQUEST FOR CONVERSION

The microorganism identified under I above was received by this International Depositary
Authority on (date of original deposit) and a request to convert
the original deposit to a deposit under the Budapest Treaty was received by it on
(date of receipt of request for conversion)


V. INTERNATIONAL DEPOSITARY AUTHORITY

Name: Russian National Collection
of Industrial Microorganisms (VKPM)
Depositary
GNIIGenetika
Address: Russia 113545 Moscow
1 Dorozhny proezd 1

Signature (s) of person(s) having the
power to represent the International

Authority or authorized official(s):
Date: Sineoky S.F.

1 Where Rule 6.4(d) applies, such date is the date on which the status of international
depositary authority was acquired



Appendix 3

page 14

Budapest treaty on the international
recognition of the deposit of microorganisms
for the purposes of patent procedure

To
State Scientific Centre of
Russian Federation
GNIIGENETIKA
Moscow 113545
1-st Dorozhny proezd 1 Russia
name and address
of depositor

INTERNATIONAL FORM
receipt in the case of an original deposit
issued pursuant to Rule 7.1 by the
INTERNATIONAL DEPOSITARY AUTHORITY
identified at the bottom of this page

I. IDENTIFICATION OF THE MICROORGANISM

Identification reference given by the DEPOSITOR	Accession number given by the INTERNATIONAL DEPOSITARY- AUTHORITY:
Escherichia coli VL2151 (pUC21)	VKPM B- 7715

II. SCIENTIFIC DESCRIPTION AND/OR PROPOSED TAXONOMIC DESIGNATION

The microorganism identified under I above was accompanied by:

☒ a scientific description Producer of proline

☐ a proposed taxonomic designation
(Mark with a cross where applicable)

III. RECEIPT AND ACCEPTANCE

This International Depositary Authority accepts the microorganism identified under I above,
1 which was received by it on 29.12.1998 (date of original deposit)

IV. RECEIPT OF REQUEST FOR CONVERSION

The microorganism identified under I above was received by this International Depositary
Authority
on (date of original deposit) and a request to convert
the
original deposit to a deposit under the Budapest Treaty was received by it on
(date of receipt of request for conversion)

V. INTERNATIONAL DEPOSITARY AUTHORITY

Name: Russian National Collection
of Industrial Microorganisms (VKPM)
Depositary
GNIIGenetika
Address: Russia 113545 Moscow
1 Dorozhny proezd 1

Signature (s) of person(s) having the
power to represent the International

Authority or authorized official(s):
Date Sineoky S.P.

1 Where Rule 6.4(d) applies, such date is the date on which the status of international
depositary authority was acquired



[Handwritten signature]

65622 E 255460

Appendix 3

page 14

Budapest treaty on the international
recognition of the deposit of microorganisms
for the purposes of patent procedure

To
State Scientific Centre of
Russian Federation
GNIIGENETIKA
Moscow 113545
1-st Dorozhny proezd 1 Russia
name and address
of depositor

INTERNATIONAL FORM
receipt in the case of an original deposit
issued pursuant to Rule 7.1 by the
INTERNATIONAL DEPOSITARY AUTHORITY
identified at the bottom of this page

I. IDENTIFICATION OF THE MICROORGANISM

Identification reference given by the DEPOSITOR	Accession number given by the INTERNATIONAL DEPOSITARY- AUTHORITY:
Escherichia coli W3350 argE::Tn10 (pKA10)(pYGGA)	VKPM B- 7716

II. SCIENTIFIC DESCRIPTION AND/OR PROPOSED TAXONOMIC DESIGNATION

The microorganism identified under I above was accompanied by:

☒ a scientific description ☐ Producer of arginine

☐ a proposed taxonomic designation
(Mark with a cross where applicable)

III. RECEIPT AND ACCEPTANCE

This International Depositary Authority accepts the microorganism identified under I above,
1 which was received by it on 29.12.1998 (date of original deposit)

IV. RECEIPT OF REQUEST FOR CONVERSION

The microorganism identified under I above was received by this International Depositary
Authority on (date of original deposit) and a request to convert
the original deposit to a deposit under the Budapest Treaty was received by it on
(date of receipt of request for conversion)



V. INTERNATIONAL DEPOSITARY AUTHORITY

Name: Russian National Collection
of Industrial Microorganisms (VKPM)
Depositary
GNIIGenetika
Address: Russia 113545 Moscow
1 Dorozhny proezd 1

Signature (s) of person(s) having the
power to represent the International

Authority or authorized official(s):
Date: Sineoky S.P.

1 Where Rule 6.4(d) applies, such date is the date on which the status of international
depositary authority was acquired



Appendix 3

page 14

Budapest treaty on the international
recognition of the deposit of microorganisms
for the purposes of patent procedure

To
State Scientific Centre of
Russian Federation
GNIIGENETIKA
Moscow 113545
1st Dorozhny proezd 1 Russia
name and address
of depositor

INTERNATIONAL FORM
receipt in the case of an original deposit
issued pursuant to Rule 7.1 by the
INTERNATIONAL DEPOSITARY AUTHORITY
identified at the bottom of this page

I. IDENTIFICATION OF THE MICROORGANISM

Identification reference given by the DEPOSITOR	Accession number given by the INTERNATIONAL DEPOSITARY- AUTHORITY:
Escherichia coli W 3350 argE::Tn10 (pKA10)(pOK12)	VKPM B- 7718

II. SCIENTIFIC DESCRIPTION AND/OR PROPOSED TAXONOMIC DESIGNATION

The microorganism identified under I above was accompanied by:

☒ a scientific description Producer of arginine

☐ a proposed taxonomic designation
(Mark with a cross where applicable)

III. RECEIPT AND ACCEPTANCE

This International Depositary Authority accepts the microorganism identified under I above,
1 which was received by it on 29.12.1998 (date of original deposit)


IV. RECEIPT OF REQUEST FOR CONVERSION

The microorganism identified under I above was received by this International Depositary
Authority
on (date of original deposit) and a request to convert
the
original deposit to a deposit under the Budapest Treaty was received by it on
(date of receipt of request for conversion)

V. INTERNATIONAL DEPOSITARY AUTHORITY

Name: Russian National Collection of Industrial Microorganisms (VKPM) GNIIGenetika Address: Russia 113545 Moscow 1 Dorozhny proezd 1	Signature (s) of person(s) having the power to represent the International Authority or authorized official(s): Date: Sineoky SP
--	---

1 Where Rule 6.4(d) applies, such date is the date on which the status of international
depository authority was acquired



To
State Scientific Centre of
Russian Federation
GNIIGENETIKA
.
Moscow 113545
!-st Dorozhny proezd 1 Russia
name and address
of depositor

INTERNATIONAL FORM
receipt in the case of an original deposit
issued pursuant to Rule 7.1 by the
INTERNATIONAL DEPOSITORY AUTHORITY
identified at the bottom of this page

Identification reference given by the
DEPOSITOR

Escherichia coli VL614 (pYGGA)

Accession number given by the
INTERNATIONAL DEPOSITARY-
AUTHORITY:

VKPM B-7719

II. SCIENTIFIC DESCRIPTION AND/OR PROPOSED TAXONOMIC DESIGNATION

The microorganism identified under I above was accompanied by:

x a scientific description

Producer of lysine

_____ a proposed taxonomic designation
(Mark with a cross where applicable)

III. RECEIPT AND ACCEPTANCE

This International Depositary Authority accepts the microorganism identified under I above,

1 which was received by it on 29.12.1998

(date of original deposit)

IV. RECEIPT OF REQUEST FOR CONVERSION

The microorganism identified under I above was received by this International Depository Authority

on _____ (date of original deposit) and a request to convert
the _____

original deposit to a deposit under the Budapest Treaty was received by it on
(date of receipt of request for conversion)

V. INTERNATIONAL DEPOSITORY AUTHORITY


Name: Russian National Collection
of Industrial Microorganisms (VKPM)
Depository
GNIIgenetika
Address: Russia 113545 Moscow
1 Dorozhny proezd 1

Signature (s) of person(s) having the power * to represent the International

Authority or authorized official(s):

Date: 12/12/2019 Sineoky, S.P.

1 Where Rule 6.4(d) applies, such date is the date on which the status of international
depository authority was acquired



Appendix 3

page 14

Budapest treaty on the international
recognition of the deposit of microorganisms
for the purposes of patent procedure

To
State Scientific Centre of
Russian Federation
GNIIGENETIKA
Moscow 113545
1-st Dorozhny proezd 1 Russia
name and address
of depositor

INTERNATIONAL FORM
receipt in the case of an original deposit
issued pursuant to Rule 7.1 by the
INTERNATIONAL DEPOSITARY AUTHORITY
identified at the bottom of this page

I. IDENTIFICATION OF THE MICROORGANISM

Identification reference given by the DEPOSITOR	Accession number given by the INTERNATIONAL DEPOSITARY- AUTHORITY:
Escherichia coli VL614 (pOK12)	VKPM E- 7722

II. SCIENTIFIC DESCRIPTION AND/OR PROPOSED TAXONOMIC DESIGNATION

The microorganism identified under I above was accompanied by:

☒ a scientific description Producer of lysine

☐ a proposed taxonomic designation
(Mark with a cross where applicable)

III. RECEIPT AND ACCEPTANCE

This International Depositary Authority accepts the microorganism identified under I above,
1 which was received by it on 29.12.1998 (date of original deposit)

IV. RECEIPT OF REQUEST FOR CONVERSION

The microorganism identified under I above was received by this International Depositary
Authority
on (date of original deposit) and a request to convert
the
original deposit to a deposit under the Budapest Treaty was received by it on
(date of receipt of request for conversion)

V. INTERNATIONAL DEPOSITARY AUTHORITY

Name: Russian National Collection
of Industrial Microorganisms (VKPM)
Depositary
GNIIGenetika
Address: Russia 113545 Moscow
1 Dorozhny proezd 1

Signature (s) of person(s) having the
power to represent the International

Authority or authorized official(s):

Date: Sineokv S.P.

1 Where Rule 6.4(d) applies, such date is the date on which the status of international
depositary authority was acquired

SECRET 6455460

Appendix 3

page 14

Budapest treaty on the international
recognition of the deposit of microorganisms
for the purposes of patent procedure

To
State Scientific Centre of
Russian Federation
GNIIGENETIKA
Moscow 113545
1-st Dorozhny proezd 1 Russia
name and address
of depositor

INTERNATIONAL FORM
receipt in the case of an original deposit
issued pursuant to Rule 7.1 by the
INTERNATIONAL DEPOSITARY AUTHORITY
identified at the bottom of this page

I. IDENTIFICATION OF THE MICROORGANISM

Identification reference given by the
DEPOSITOR

Escherichia coli AJ13199(pUC21)

Accession number given by the
INTERNATIONAL DEPOSITARY-
AUTHORITY:
VKPM B- 7728

II. SCIENTIFIC DESCRIPTION AND/OR PROPOSED TAXONOMIC DESIGNATION

The microorganism identified under I above was accompanied by:

☒ a scientific description Producer of glutamic acid

☐ a proposed taxonomic designation
(Mark with a cross where applicable)

III. RECEIPT AND ACCEPTANCE

This International Depositary Authority accepts the microorganism identified under I above,
1 which was received by it on 29.12.1998 (date of original deposit)

IV. RECEIPT OF REQUEST FOR CONVERSION

The microorganism identified under I above was received by this International Depositary
Authority
on (date of original deposit) and a request to convert
the
original deposit to a deposit under the Budapest Treaty was received by it on
(date of receipt of request for conversion)

V. INTERNATIONAL DEPOSITARY AUTHORITY

Name: Russian National Collection
of Industrial Microorganisms (VKPM)
Depositary
GNIIGenetika
Address: Russia 113545 Moscow
1 Dorozhny proezd 1

Signature (s) of person(s) having the
power to represent the International

Authority or authorized official(s):
Date: Sineoky S.P.

1 Where Rule 6.4(d) applies, such date is the date on which the status of international
depositary authority was acquired

Appendix 3

page 14

Budapest treaty on the international
recognition of the deposit of microorganisms
for the purposes of patent procedure

To
State Scientific Centre of
Russian Federation
GNIIGENETIKA
Moscow 113545
1-st Dorozhny proezd 1 Russia
name and address
of depositor

INTERNATIONAL FORM
receipt in the case of an original deposit
issued pursuant to Rule 7.1 by the
INTERNATIONAL DEPOSITARY AUTHORITY
identified at the bottom of this page

I. IDENTIFICATION OF THE MICROORGANISM

Identification reference given by the
DEPOSITOR

Escherichia coli AJ13199(pYAHN)

Accession number given by the
INTERNATIONAL DEPOSITARY-
AUTHORITY:

VKPM B- 7729

II. SCIENTIFIC DESCRIPTION AND/OR PROPOSED TAXONOMIC DESIGNATION

The microorganism identified under I above was accompanied by:

☒ a scientific description

Producer of glutamic acid

☐ a proposed taxonomic designation
(Mark with a cross where applicable)

III. RECEIPT AND ACCEPTANCE

This International Depositary Authority accepts the microorganism identified under I above,

1

which was received by it on 29.12.1998

(date of original deposit)

IV. RECEIPT OF REQUEST FOR CONVERSION

The microorganism identified under I above was received by this International Depositary
Authority

on


(date of original deposit) and a request to convert

the

original deposit to a deposit under the Budapest Treaty was received by it on
(date of receipt of request for conversion)

V. INTERNATIONAL DEPOSITARY AUTHORITY

Name: Russian National Collection
of Industrial Microorganisms (VKPM)
GNIIGenetika
Address: Russia 113545 Moscow
1 Dorozhny proezd 1



Signature (s) of person(s) having the
power to represent the International

Authority or authorized official(s):
Date: Sineoky S.P.

1 Where Rule 6.4(d) applies, such date is the date on which the status of international
depositary authority was acquired

Appendix 3

page 14

Budapest treaty on the international
recognition of the deposit of microorganisms
for the purposes of patent procedure

To
State Scientific Centre of
Russian Federation
GNIIGENETIKA
Moscow 113545
1-st Dorozhny proezd 1 Russia
name and address
of depositor

INTERNATIONAL FORM
receipt in the case of an original deposit
issued pursuant to Rule 7.1 by the
INTERNATIONAL DEPOSITARY AUTHORITY
identified at the bottom of this page

I. IDENTIFICATION OF THE MICROORGANISM

Identification reference given by the
DEPOSITOR

Escherichia coli AJ13199(pYFIK)

Accession number given by the
INTERNATIONAL DEPOSITARY-
AUTHORITY:
VKPM B- 7730

II. SCIENTIFIC DESCRIPTION AND/OR PROPOSED TAXONOMIC DESIGNATION

The microorganism identified under I above was accompanied by:

☒ a scientific description Producer of glutamic acid
☐ a proposed taxonomic designation
(Mark with a cross where applicable)

III. RECEIPT AND ACCEPTANCE

This International Depositary Authority accepts the microorganism identified under I above,
1 which was received by it on 29.12.1998 (date of original deposit)

IV. RECEIPT OF REQUEST FOR CONVERSION

The microorganism identified under I above was received by this International Depositary
Authority
on (date of original deposit) and a request to convert
the
original deposit to a deposit under the Budapest Treaty was received by it on
(date of receipt of request for conversion)


V. INTERNATIONAL DEPOSITARY AUTHORITY

Name: Russian National Collection
of Industrial Microorganisms (VKPM)
Depositary
GNIIGenetika
Address: Russia 113545 Moscow
1 Dorozhny proezd 1

Signature(s) of person(s) having the
power to represent the International

Authority or authorized official(s):
Date: Sineoky S.P.

1 Where Rule 6.4(d) applies, such date is the date on which the status of international
depositary authority was acquired



04455450

Appendix 3
page 14

Budapest treaty on the international
recognition of the deposit of microorganisms
for the purposes of patent procedure

To
State Scientific Centre of
Russian Federation
GNIIGENETIKA
Moscow 113545
1-st Dorozhny proezd 1 Russia
name and address
of depositor

INTERNATIONAL FORM
receipt in the case of an original deposit
issued pursuant to Rule 7.1 by the
INTERNATIONAL DEPOSITARY AUTHORITY
identified at the bottom of this page

I. IDENTIFICATION OF THE MICROORGANISM

Identification reference given by the DEPOSITOR	Accession number given by the INTERNATIONAL DEPOSITARY- AUTHORITY:
Escherichia coli AJ13199(pYEAS)	VKPM B- 7731

II. SCIENTIFIC DESCRIPTION AND/OR PROPOSED TAXONOMIC DESIGNATION

The microorganism identified under I above was accompanied by:

☒ a scientific description Producer of glutamic acid
☐ a proposed taxonomic designation
(Mark with a cross where applicable)

III. RECEIPT AND ACCEPTANCE

This International Depositary Authority accepts the microorganism identified under I above,
I which was received by it on 29.12.1998 (date of original deposit)

IV. RECEIPT OF REQUEST FOR CONVERSION

The microorganism identified under I above was received by this International Depositary
Authority on (date of original deposit) and a request to convert
the original deposit to a deposit under the Budapest Treaty was received by it on
(date of receipt of request for conversion)


V. INTERNATIONAL DEPOSITARY AUTHORITY

Name: Russian National Collection
of Industrial Microorganisms (VKPM)
Depositary
GNIIGenetika
Address: Russia 113545 Moscow
1 Dorozhny proezd 1

Signature (s) of person(s) having the
power to represent the International

Authority or authorized official(s):
Date: Sineoky S.P.

1 Where Rule 6.4(d) applies, such date is the date on which the status of international
depositary authority was acquired



Appendix 3
page 14

Budapest treaty on the international
recognition of the deposit of microorganisms
for the purposes of patent procedure

To
State Scientific Centre of
Russian Federation
GNIIGENETIKA
Moscow 113545
1-st Dorozhny proezd 1 Russia
name and address
of depositor

INTERNATIONAL FORM
receipt in the case of an original deposit
issued pursuant to Rule 7.1 by the
INTERNATIONAL DEPOSITARY AUTHORITY
identified at the bottom of this page

I. IDENTIFICATION OF THE MICROORGANISM

Identification reference given by the
DEPOSITOR

Escherichia coli VL2151 (pYAHN)

Accession number given by the
INTERNATIONAL DEPOSITARY-
AUTHORITY:
VKPM B- 7748

II. SCIENTIFIC DESCRIPTION AND/OR PROPOSED TAXONOMIC DESIGNATION

The microorganism identified under I above was accompanied by:

☒ a scientific description Producer of proline

☐ a proposed taxonomic designation
(Mark with a cross where applicable)

III. RECEIPT AND ACCEPTANCE

This International Depositary Authority accepts the microorganism identified under I above,
1 which was received by it on 29.12.1998 (date of original deposit)

IV. RECEIPT OF REQUEST FOR CONVERSION

The microorganism identified under I above was received by this International Depositary
Authority
on (date of original deposit) and a request to convert
the
original deposit to a deposit under the Budapest Treaty was received by it on
(date of receipt of request for conversion)

V. INTERNATIONAL DEPOSITARY AUTHORITY

Name: Russian National Collection
of Industrial Microorganisms (VKPM)
Depositary
GNIIGenetika
Address: Russia 113545 Moscow
1 Dorozhny proezd 1

Signature (s) of person(s) having the
power to represent the International

Authority or authorized official(s):
Date: Sineoky S.P.

1 Where Rule 6.4(d) applies, such date is the date on which the status of international
depositary authority was acquired

Appendix 3
page 14
Budapest treaty on the international
recognition of the deposit of microorganisms
for the purposes of patent procedure

To
State Scientific Centre of
Russian Federation
GNIIGENETIKA
Moscow 113545
1-st Dorozhny proezd 1 Russia
name and address
of depositor

INTERNATIONAL FORM
receipt in the case of an original deposit
issued pursuant to Rule 7.1 by the
INTERNATIONAL DEPOSITARY AUTHORITY
identified at the bottom of this page

I. IDENTIFICATION OF THE MICROORGANISM

Identification reference given by the DEPOSITOR	Accession number given by the INTERNATIONAL DEPOSITARY- AUTHORITY:
Escherichia coli VL2160 (pUC21)	VKPM B- 7752

II. SCIENTIFIC DESCRIPTION AND/OR PROPOSED TAXONOMIC DESIGNATION

The microorganism identified under I above was accompanied by:

☒ a scientific description Producer of histidine

☐ a proposed taxonomic designation
(Mark with a cross where applicable)

III. RECEIPT AND ACCEPTANCE

This International Depositary Authority accepts the microorganism identified under I above,
1 which was received by it on 29.12.1998 (date of original deposit)

IV. RECEIPT OF REQUEST FOR CONVERSION

The microorganism identified under I above was received by this International Depositary
Authority
on (date of original deposit) and a request to convert
the
original deposit to a deposit under the Budapest Treaty was received by it on
(date of receipt of request for conversion)



V. INTERNATIONAL DEPOSITARY AUTHORITY

Name: Russian National Collection
of Industrial Microorganisms (VKPM)
Depositary
GNIIGenetika
Address: Russia 113545 Moscow
1 Dorozhny proezd 1

Signature (s) of person(s) having the
power to represent the International

authority or authorized official(s):
Date: Sineoky SP

1 Where Rule 6.4(d) applies, such date is the date on which the status of international
depositary authority was acquired



Appendix 3

page 14

Budapest treaty on the international
recognition of the deposit of microorganisms
for the purposes of patent procedure

To
State Scientific Centre of
Russian Federation
GNIIGENETIKA
Moscow 113545
1-st Dorozhny proezd 1 Russia
name and address
of depositor

INTERNATIONAL FORM
receipt in the case of an original deposit
issued pursuant to Rule 7.1 by the
INTERNATIONAL DEPOSITARY AUTHORITY
identified at the bottom of this page

I. IDENTIFICATION OF THE MICROORGANISM

Identification reference given by the
DEPOSITOR

Escherichia coli VL2160 (pYEAS)

Accession number given by the
INTERNATIONAL DEPOSITARY-
AUTHORITY:
VKPM B- 7753

II. SCIENTIFIC DESCRIPTION AND/OR PROPOSED TAXONOMIC DESIGNATION

The microorganism identified under I above was accompanied by:

☒ a scientific description Producer of histidine

☐ a proposed taxonomic designation
(Mark with a cross where applicable)

III. RECEIPT AND ACCEPTANCE

This International Depositary Authority accepts the microorganism identified under I above,
1 which was received by it on 29.12.1998 (date of original deposit)

IV. RECEIPT OF REQUEST FOR CONVERSION

The microorganism identified under I above was received by this International Depositary
Authority
on (date of original deposit) and a request to convert
the
original deposit to a deposit under the Budapest Treaty was received by it on
(date of receipt of request for conversion)

V. INTERNATIONAL DEPOSITARY AUTHORITY

Name: Russian National Collection
of Industrial Microorganisms (VKPM)
Depositary
GNIIGenetika
Address: Russia 113545 Moscow
1 Dorozhny proezd 1

Signature (s) of person(s) having the
power to represent the International

Authority or authorized official(s):

Date:

Sineoky S.P.

1 Where Rule 6.4(d) applies, such date is the date on which the status of international
depositary authority was acquired

Appendix 3

page 14

Budapest treaty on the international
recognition of the deposit of microorganisms
for the purposes of patent procedure

To
State Scientific Centre of
Russian Federation
GNIIGENETIKA
Moscow 113545
1-st Dorozhny proezd 1 Russia
name and address
of depositor

INTERNATIONAL FORM
receipt in the case of an original deposit
issued pursuant to Rule 7.1 by the
INTERNATIONAL DEPOSITARY AUTHORITY
identified at the bottom of this page

I. IDENTIFICATION OF THE MICROORGANISM

Identification reference given by the DEPOSITOR	Accession number given by the INTERNATIONAL DEPOSITARY- AUTHORITY:
Escherichia coli VL2160 (pYFIK)	VKPM B- 7754

II. SCIENTIFIC DESCRIPTION AND/OR PROPOSED TAXONOMIC DESIGNATION

The microorganism identified under I above was accompanied by:

☒ a scientific description Producer of histidine☐ a proposed taxonomic designation
(Mark with a cross where applicable)

III. RECEIPT AND ACCEPTANCE

This International Depositary Authority accepts the microorganism identified under I above,
1 which was received by it on 29.12.1998 (date of original deposit)

IV. RECEIPT OF REQUEST FOR CONVERSION

The microorganism identified under I above was received by this International Depositary
Authority
on (date of original deposit) and a request to convert
the
original deposit to a deposit under the Budapest Treaty was received by it on
(date of receipt of request for conversion)

V. INTERNATIONAL DEPOSITARY AUTHORITY

Name: Russian National Collection
of Industrial Microorganisms (VKPM)
Depositary
GNIIGenetika
Address: Russia 113545 Moscow
1 Dorozhny proezd 1

Signature (s) of person(s) having the
power to represent the International

Authority or authorized official(s):
Date: Sineoky S.P.

1 Where Rule 6.4(d) applies, such date is the date on which the status of international
depositary authority was acquired

